MARK SCHEME for the October/November 2011 question paper

for the guidance of teachers

9700 BIOLOGY

9700/42

Paper 4 (A2 Structured Questions), maximum raw mark 100

MMM. Hiremepapers.com

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2011 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



| Page 2 | Mark Scheme: Teachers' version | Syllabus | Paper |
|--------|----------------------------------------|----------|-------|
| | GCE AS/A LEVEL – October/November 2011 | 9700 | 42 |

Mark scheme abbreviations:

- ; separates marking points
- *I* alternative answers for the same point
- R reject
- A accept (for answers correctly cued by the question, or by extra guidance)
- **AW** alternative wording (where responses vary more than usual)
- **<u>underline</u>** actual word given must be used by candidate (grammatical variants excepted)
- max indicates the maximum number of marks that can be given
- ora or reverse argument
- mp marking point (with relevant number)
- ecf error carried forward
- I ignore
- **AVP** Alternative valid point (examples given as guidance)

| | Pa | Page 3 Mark Scheme: Teachers' version S GCE AS/A LEVEL – October/November 2011 | | Syllabus | Paper | |
|---|-----|-----------------------------------------------------------------------------------|-------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------|-----------------------|
| | | | | GCE AS/A LEVEL – October/November 2011 | 9700 | 42 |
| 1 | (a) | <u>allo</u> | patric | 2; | | [1] |
| | (b) | 1. 2. 3. 4. | inbre little | ks / populations, isolated from each other ; eeding / no interbreeding ; mutation ; ' ; e.g. small population to start with / small gene pool t | o start with | [2 max] |
| | (c) | 1. 2. 3. 4. | <i>idea</i> hunt | culture / buildings / AW ; of wolves dying ; ing / trapping / AW ; idisation / infertility / change in (wolf) gene pool / loss o | of wolf alleles / A | W; [4] |
| | (d) | allo | %) ;; w on rking | e mark for number not rounded up or incorrect answ | ver but correct | [2] idea regarding |
| | | | | | | [Total: 9] |
| 2 | (a) | 1. 2. | | three colours (for positive reactions) / only a small ran neasurement of actual concentration / no numerical va | | [2] |
| | (b) | (i) | perc | oxidase; | | [1] |
| | | (ii) | 2. 3. | (catalyses breakdown of hydrogen peroxide) to produc chromogen, oxidised by / reacts with, (oxygen) ; produces range of colours ; more, peroxide / oxygen produced, = greater change / | | [2 max] |
| | | (iii) | 2. 3. | to keep out, proteins / enzymes / polymer / named larg R large molecules unqualified to prevent interference (to reactions) ; to prevent loss of, enzyme / chromogen ; so still allowing reaction to occur ; | ge molecule ; | [2 max] |
| | (c) | (i) | 2. in A | B has diabetes and A does not; A's, values / peak, lower because he secretes insulin or B's, values / peak, higher because, no / little, insulin; (insulin affects), liver / muscle, cells; | | |
| | | | 4. 5. 6. <i>in B</i> | increase in glucose uptake / increase in permeability c increase in use of glucose in respiration ; (more) glucose converted to glycogen ; | f membranes (to | o glucose) ; |
| | | | | because cells unresponsive to insulin ; ept quoted values for lower and higher in mark point 2 | | [4 max] |

| Page 4 | Mark Scheme: Teachers' version | Syllabus | Paper |
|--------|----------------------------------------|----------|-------|
| | GCE AS/A LEVEL – October/November 2011 | 9700 | 42 |

- (ii) (the concentration of blood glucose), above which some glucose appears in the urine / AW;
- (iii) 1. (at first), glucose reabsorbed by proximal convoluted tubule ;
 - 2. ref. co transported with Na^+ / facilitated diffusion / protein carrier ;
 - 3. above 180mg (100cm⁻³ glucose in blood) no further reabsorption ;
 - 4. because carriers (in PCT) saturated / AW ;

[Total: 15]

[3 max]

3 (a)

| | male | | female | |
|---|-------------------------------------------|----|----------------------------------|---|
| 1 | produces sperm or produces, oocyte | | produces, oocyte | ; |
| 2 | division of cytoplasm is equal | or | division of cytoplasm is unequal | ; |
| 3 | four gametes produced | or | one gamete produced | ; |
| 4 | no polar bodies | or | polar bodies | ; |
| 5 | ref. maturation | or | no equivalent maturation stage | ; |
| 6 | ref. meiosis completed | or | ref. incomplete meiosis | ; |

[3 max]

[2 max]

- (b) 1. a ductless gland;
 - 2. hormones in the blood;
 - 3. ref. target, organ / tissues;
- (c) 1. (both), reduce / stop, secretion (of FSH and LH);
 - 2. (both) involve negative feedback;
 - 3. to, anterior pituitary / hypothalamus;
 - 4. both are, contraceptives / description ;

[3 max]

[Total: 8]

- **4** (a) 1. low oxygen (in water) results in anaerobic respiration ;
 - 2. (anaerobic respiration) produces alcohol;
 - 3. rice tolerant to alcohol;
 - 4. (because rice has) high levels of, alcohol dehydrogenase / enzyme that breaks down alcohol;
 - 5. presence of, aerenchyma / described ;
 - 6. allows, oxygen / air, to reach roots (from aerial tissues); [3 max]
 - (b) (i) 1. (immersion in water) stimulates production of ethene;
 - 2. (concentration of) ethene produced increased with time (after submergence);
 - 3. very little difference in ethene production between T65 and C9285;
 - 4. use of figures ; 2 values of ethene **plus** 2 values of time for **either** T65 **or** C9285

[2 max]

| Page 5 | | | Mark Scheme: Teachers' version | Syllabus | Paper | |
|--------|-----|----------------------------------------|----------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------|-------------------------------------------|
| | | | | GCE AS/A LEVEL – October/November 2011 | 9700 | 42 |
| | | (ii) | 2. 3. | in T65 ethene does not affect internode elongation b internode elongation ; in C9285, greater concentrations of ethene cause grea use of comparative figures to support mark point 1 or <i>least once</i> | ater elongation ; | bene promotes both units at [2 max] |
| | (c) | 1. 2. 3. 4. | incr GA s | genes present in C9285 / <i>SK</i> genes not present in T65 eased production of GA in C9285 / little or no increased p stimulates, stem elongation / AW ; ; e.g. T65 has no receptors for ethene | | in T65 ; [3 max] |
| | (d) | (i) | 0. n or | more important; ora <i>ivara</i> has mutated <i>SK2</i> and does not have deepwater i <i>lumaepatula</i> has <i>SK2</i> but not <i>SK1</i> and does have deep | · | ; [2] |
| | | (ii) | 1. 2. | (addition / insertion), of a, base / nucleotide, to DNA / t changes a, sequence of three bases / triplet / codon ; (triplet) no longer codes for an amino acid ; | o a gene ; | |
| | | (iii) | 2. 3. | breed deepwater variety with (high-yielding) non-deep identify / select, offspring with both deepwater respons breed selected offspring (with both deepwater respons continue for many generations ; | se and high yield | |
| | | | | | | [Total: 17] |
| 5 | (a) | 1. 2. 3. 4. 5. 6. 7. | caus deliv only ease serio | eed by a single gene ; sed by a recessive allele ; rery of, correct / dominant / normal, allele (could correc need to get allele into a few cells ; e of access to affected area ; bus so worth the risk ; ; e.g. only targets eye / no surgery needed | t the condition) | [3 max] |
| | (b) | 1. 2. 3. | corre | s no longer able to cause infections ; ect / dominant / normal, allele (of <i>RPE65</i>) added ; noter added ; | | [2 max] |
| | (c) | 2. 3. | rare expe | to safety / not known if the technique might have side e condition ; ense ; ' ; e.g. trial to see if delivery method works | effects; | [2 max] |
| | | | | | | [Total: 7] |
| 6 | (a) | (i) | phos | sphorylation; | | [1] |
| | . / | | lysis | | | [1] |
| | | . / | | | | |

| Page | 6 | Mark Scheme: Teachers' version | Syllabus | Paper |
|--------------------------|-----------------------|---------------------------------------------------------------------------------------------|----------|---------|
| | | GCE AS/A LEVEL – October/November 2011 | 9700 | 42 |
| (iii) | dehy | /drogenation / oxidation; ignore reduction of NAD | | [1] |
| | | activation energy / AW ; plit / AW ; | | [2] |
| (c) 1. 2. 3. 4. | etha etha by re | arboxylated / carbon dioxide given off ; nal produced ; nal reduced ; educed NAD ; | | |
| 5. 6. | | hanol ; /drogenase ; | | [4 max] |

[Total: 9]

| _ |
|---|
| 7 |
| 1 |

| step | reason for step |
|------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------|
| obtain copies of gene with sticky ends | the gene codes for the synthesis of insulin |
| plasmid (used) ; | acts as a vector for the transfer of the gene into the host |
| use restriction endonuclease enzyme | to produce 'sticky ends' or cut at specific, site / sequence ; |
| mix vector and gene | gene inserts into, vector / plasmid or forms recombinant DNA / AW ; |
| | A detail of complementary base pairing |
| (use DNA) ligase ; | to seal the sugar-phosphate backbone |
| insert, plasmid / vector, into host / <i>E. coli / bacteria</i> ; | to obtain transformed host <i>E. coli</i> cells |
| screen for, and obtain, successfully transformed cells | so only recombinant host cells cultured / AW; |
| ref. batch / continuous, culture or fermenter or bacterial cloning / population growth ; | to obtain large amounts of insulin for extraction and purification |

[Total: 7]

| | Page | 7 | М | ark Sche | eme: Teac | hers' ve | rsion | | Sy | llabus | Pa | aper |
|---|----------------------|--------------------|---------------------------------------------------------------|----------|------------|----------|--------|---------|---------|-----------|-------|---------|
| | | | GCE AS | 6/A LEVE | EL – Octob | er/Nove | ember | 2011 | g | 700 | | 42 |
| 8 | (c na ai pe | amed a nimal fu | e) smoke ; ir pollutant ır / skin flak s / aerosol / | es / AW | | | | | | | | |
| | | | spores ; | | | | | | | | | [2 max] |
| | (b) (i |) high | tensile str | ength / | withstands | pulling | forces | / fibro | us / in | soluble / | forms | fibri |

[1]

(ii) 1. inbreeding;

flexible;

- 2. little genetic variation / small gene pool / small population ;
- 3. many carried faulty allele / AW;
- 4. faulty / mutant, allele, could be dominant or recessive ;
- 5. little phenotypic variation ;

[3 max]

[Total: 6]

9 (a) dominant

(allele) that always expresses itself (in the phenotype) when present **or**

(allele) which influences the phenotype even in the presence of an alternative allele ;

gene

length of DNA / sequence of nucleotides, coding for a (specific) polypeptide; A protein [2]

(b)

| parental phenotypes | man without TSC | woman with TSC | | |
|---------------------------------|----------------------|----------------|---|-----|
| parental genotypes | tt | Tt | | |
| gametes | all t | T or t | ; | |
| offspring genotypes | Tt | tt | | |
| offspring phenotypes | TSC | normal | ; | |
| probability of child having TSC | 50% / 0.50 / 1in 2 ; | | | |
| | | | | [3] |

- (c) 1. spontaneous / random / chance ;
 - 2. mutation of, gene / allele;
 - 3. AVP ; e.g. named mutagen / detail of mutation

[2 max]

[Total: 7]

| Page 8 | Mark Scheme: Teachers' version | Syllabus | Paper |
|--------|----------------------------------------|----------|-------|
| | GCE AS/A LEVEL – October/November 2011 | 9700 | 42 |

- **10 (a)** 1. ground substance / stroma;
 - 2. for, light independent stage / Calvin cycle ;
 - 3. contains enzymes / named enzyme e.g. rubisco ;
 - 4. also, sugars / lipids / starch / ribosomes / DNA;
 - 5. internal membrane system;
 - 6. for, light dependent stage;
 - 7. fluid-filled sacs / thylakoids;
 - 8. grana are stacks of thylakoids;
 - 9. (grana) hold (photosynthetic) pigments ;
 - 10. (grana) have large surface area for (maximum) light absorption;
 - 11. (pigments are arranged in), light harvesting clusters / photosystems;
 - 12. primary pigment / reaction centre / chlorophyll a, surrounded by accessory pigments ;
 - 13. (accessory pigments) pass energy to, primary pigment / reaction centre / chlorophyll a ;
 - 14. different photosystems absorb light at different wavelengths ;
 - 15. membranes hold, ATP synthase / electron carriers;
 - 16. for, photophosphorylation / chemiosmosis;

[9 max]

- (b) 17. grind leaf with solvent;
 - 18. example of solvent ; e.g. propanone
 - 19. leaf extract contains mixture of pigments;
 - 20. ref. concentrate extract;
 - 21. further detail ; e.g. pencil line drawn / extract placed on chromatography paper / repetitive spotting / drying between spots
 - 22. paper placed (vertically) in jar of (different) solvent ;
 - 23. solvent rises up paper;
 - 24. each pigment travels at different speed ;
 - 25. pigments separated as they ascend;
 - 26. distance moved by each pigment is unique;
 - 27. Rf value;
 - 28. two dimensional chromatography;
 - 29. better separation of pigments ;

[6 max]

[Total: 15]

- **11** (a) 1. axon phospholipid bilayer impermeable to K^+ / Na⁺;
 - 2. sodium potassium pump;
 - 3. detail of sodium-potassium pump; e.g. transmembrane / globular / ATP binding site
 - 4. active process / ATP used / energy needed ;
 - 5. 3 Na⁺ (pumped) out / 2 K⁺ (pumped) in ;
 - 6. K⁺ diffuse out / Na⁺ diffuse in ;
 - 7. through, protein channels transport proteins ;
 - 8. more K^+ channels open than Na⁺ channels ;
 - 9. therefore, membrane more permeable to K^+ or more K^+ leave than Na⁺ enter (axon);
 - 10. inside relatively more negative than outside;
 - 11. –65mV ; **A** –70mV
 - 12. idea of leaking K⁺ responsible for resting potential / AW ;
 - 13. electrochemical gradient;
 - 14. voltage-gated channels closed;

[9 max]

| Page 9 | Mark Scheme: Teachers' version | Syllabus | Paper |
|--------|----------------------------------------|----------|-------|
| | GCE AS/A LEVEL – October/November 2011 | 9700 | 42 |

(b) general

- 15. respond to stimuli / AW;
- 16. (some) receptors are the ends of sensory neurones ;
- 17. (some) receptors are cells;
- 18. they are energy transducers;
- 19. stimulus causes sodium ion channels to open ;
- 20. sodium ions enter cell;
- 21. depolarisation;
- 22. receptor / generator, potential;
- 23. if (receptor potential) greater than threshold then action potential generated / all or nothing principle described;
- 24. increased stimulus strength leads to increased frequency of action potentials ;

examples – allow any two below

| receptor | form of energy detected |
|-----------------------------------|--------------------------|
| rods / cones | light ; |
| taste buds / olfactory cells | chemical ; |
| Pacinian \ Meissner's, corpuscle | pressure / touch ; |
| Ruffinis endings | heat ; |
| proprioreceptors | mechanical displacement; |
| hair cells in semicircular canals | movement; |
| hairs cells in cochlea | sound; |

[max 2] [6 max]

[Total: 15]