

Cambridge Assessment International Education

Cambridge International Advanced Subsidiary and Advanced Level

CHEMISTRY 9701/22

Paper 2 AS Level Structured Questions

May/June 2019

MARK SCHEME
Maximum Mark: 60

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2019 series for most Cambridge IGCSE™, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.



PUBLISHED

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always whole marks (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- · marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

© UCLES 2019 Page 2 of 12

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.



© UCLES 2019 Page 3 of 12

Question	Answer	Marks			
1(a)(i)	C ₄ H ₁₀ / same molecular formula /	1			
	OR				
	same number of carbon (atoms) and hydrogen (atoms)				
	different structural formula				
	OR				
	description of different structural formula which does not imply stereoisomerism				
1(a)(ii)	structural / chain	1			
1(b)	(forward reaction is) exothermic reaction	1			
	the proportion of methylpropane / product decreases	1			
	OR				
	the proportion of butane / reactant increases				

© UCLES 2019 Page 4 of 12

Question	Answer	Marks
1(c)(i)	t shown on graph which corresponds to start of the horizontal part of both curves.	1
1(c)(ii)	concentration of butane = 0.3 mol dm ⁻³ AND concentration of methylpropane = 0.7 mol dm ⁻³	1
1(c)(iii)	[methylpropane] / [butane]	1

© UCLES 2019 Page 5 of 12

Question	Answer	Marks
1(c)(iv)	1 value for K _c	
	$K_c = \frac{\text{value of methylpropane in (ii)}}{\text{value of butane in (ii)}} = 0.7 / 0.3 = 2.3 (3)$	
	M2 units consistent with expression used in M1	
	no units / dimensionless / none	

Question	Answer	Marks			
2(a)	trend in volatility down the group	1			
	decrease (in volatility)				
	identification of specific IMF increasing				
	increasing (strength of) induced dipole (id) (interactions between molecules)				
	explanation in terms of electrons				
	increasing number of electrons				
2(b)(i)	Conditions for reaction with Cl ₂ at room temperature	1			
	ultra-violet / uv				

© UCLES 2019 Page 6 of 12

Question	Answer	Marks				
2(b)(ii)	$I_2(g/s) + H_2(g) \rightleftharpoons 2HI(g)$	1				
	1 correctly balanced equation					
	M2 correct state symbols AND use of equilibrium sign					
2(c)(i)	proton / H ⁺ donor					
2(c)(ii)	acid HC <i>l</i> AND conjugate base C <i>l</i> -					
2(c)(iii)	co-ordinate / dative (covalent)					
2(c)(iv)	(triangular / trigonal) pyramid(al)	1				
	107 ⁽⁰⁾	1				



© UCLES 2019 Page 7 of 12

Question					Answer	Marks
3(a)(i)	$SiCl_4(l) + 2H_2O(l) \rightarrow SiC$ (state symbols required)) ₂ (s) + 4H	IC <i>l</i> (aq	/g)		1
3(a)(ii)	hydrolysis					1
3(a)(iii)	NaCl – ionic					1
	SiC4 – covalent					1
3(a)(iv)	M1 statement correctly co	mparing	the diff	erence i	n electronegativity between Si and C <i>l</i> AND Na and C <i>l</i>	1
	OR					
	Na is less electronegative that Si ORA					
	M2 NaCl transfer of electr	ons				1
	M3 SiCl ₄ shared (pair of) electrons					1
3(b)(i)	Chlorine containing species	Cl ₂	HC1	HOC1		2
	Oxidation number of chlorine	0	-1	(+)1		
	Award 2 marks for 3 correct oxidation numbers					
	Award 1 Mark for 2 correct oxidation numbers					
3(b)(ii)	disproportionation				1	
3(b)(iii)	kills micro-organisms					1

© UCLES 2019 Page 8 of 12

Question	Answer	Marks
3(c)	$2NaOH + HCl + HClO \rightarrow NaCl + NaClO + 2H_2O$	2
	M1 Identifies the product NaClO	
	2 Correctly balances the equation	
	OR	
	The overall equation may be seen as: NaOH + HC l \rightarrow NaC l + H $_2$ O AND NaOH + HC l \rightarrow NaC l + H $_2$ O	

Question	Answer	Marks
4(a)	name of source crude oil / petroleum	1
	outline of separation of hydrocarbons (separation of molecules according to) different boiling points	1
4(b)(i)	cracking	1
4(b)(ii)	$2C_7H_{16} \rightarrow C_4H_{10} + C_6H_{14} + 2C_2H_4$	1

© UCLES 2019 Page 9 of 12

Question		Ansv	ver	Marks	
4(c)		method 1	method 2		
	M1	1 / 28 (= 0.035714)	1:88/28 (= 3.14286)	1	
	M2	2 × M1 (= 1 / 14 = 0.07143)	M1 / 44 (= 0.071429)	1	
	M3	$M2 \times 24 = 1.7 \text{ dm}^3$	$M2 \times 24 = 1.7 \text{ dm}^3$	1	
4(d)(i)	unpaired electro	on(s)		1	
4(d)(ii)	homolytic (fissio	n)		1	
	one electron goes to each chlorine / atom				
	OR				
	pair of electrons	is shared out (equally between the two atoms)			
4(d)(iii)	propagation			1	
4(d)(iv)	CH ₄ + (•)C <i>l</i> →	(•)CH ₃ + HC l OR CH ₄ + (•)C l \rightarrow HC l + CH ₃ (•	•)	1	

© UCLES 2019 Page 10 of 12

Question	Answer	Marks		
4(d)(v)	M1 identify bond types in alkanes and alkenes	1		
	bonds in alkanes are (all) sigma / σ			
	AND			
	alkenes have (C = C made of) sigma AND pi / σ AND π			
	M2 electrons in π (of the C = C) are responsible for the reaction			
	electrophiles are attracted to / attack to electrons in $\operatorname{\mathbf{pi}}$ / π			
	OR			
	electrophiles react with pi / π			

Question	Answer	Marks
5(a)(i)	-)propyl ethanoate	
5(a)(ii)	aOH / sodium hydroxide	
5a(iii)	H H H H-C-C-O, H H H	1
5(a)(iv)	perfume / solvent	1

© UCLES 2019 Page 11 of 12

Question			Answer		Marks
5(b)(i)	M1 divide by A _r	С	Н	0	1
		54.5 / 12	9.1/1	36.4 / 16	
	M2 divide by smallest i	number			1
		4.54 / 2.275 (= 2 OR 1.99)	9.1 / 2.275(=4)	2.275 / 2.275 (=1)	
	M3 empirical formula b	ased on correctly rounded value	s of M2		1
			C_2H_4O		
5(b)(ii)	(relative) molecular ma	ss/M _r			1
5(c)(i)	C ₃ H ₆ O ₂				1
5(c)(ii)	X and Z – no reaction /	no (visible) change			1
	Y – effervesces				1
5(d)	2HCO ₂ H + Na ₂ CO ₃ →	2HCO ₂ Na + CO ₂ + H ₂ O			1



© UCLES 2019 Page 12 of 12