

OXFORD

INTERNATIONAL
AQA EXAMINATIONS

INTERNATIONAL AS Chemistry

(9620)

CH02: Organic 1 and Physical 1
Mark Scheme

June 2017

Version: 1.0 Final

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Assessment Writer.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

A-level Chemistry

Mark Scheme Instructions for Examiners

1. General

The mark scheme for each question shows:

- the marks available for each part of the question
- the total marks available for the question
- the typical answer or answers which are expected
- extra information to help the examiner make his or her judgement and help to delineate what is acceptable or not worthy of credit or, in discursive answers, to give an overview of the area in which a mark or marks may be awarded.

The extra information in the 'Comments' column is aligned to the appropriate answer in the left-hand part of the mark scheme and should only be applied to that item in the mark scheme.

You should mark according to the contents of the mark scheme. If you are in any doubt about applying the mark scheme to a particular response, consult your Team Leader.

At the beginning of a part of a question a reminder may be given, for example: where consequential marking needs to be considered in a calculation; or the answer may be on the diagram or at a different place on the script.

In general the right-hand side of the mark scheme is there to provide those extra details which might confuse the main part of the mark scheme yet may be helpful in ensuring that marking is straightforward and consistent.

The use of M1, M2, M3 etc in the right-hand column refers to the marking points in the order in which they appear in the mark scheme. So, M1 refers to the first marking point, M2 the second marking point etc.

2. Boldening

- 2.1** In a list of acceptable answers where more than one mark is available 'any **two** from' is used, with the number of marks emboldened. Each of the following bullet points is a potential mark.
- 2.2** A bold **and** is used to indicate that both parts of the answer are required to award the mark.
- 2.3** Alternative answers acceptable for a mark are indicated by the use of **OR**. Different terms in the mark scheme are shown by a / ; eg allow smooth / free movement.

3. Marking points

3.1 Marking of lists

This applies to questions requiring a set number of responses, but for which students have provided extra responses. The general 'List' principle to be followed in such a situation is that 'right + wrong = wrong'.

Each error / contradiction negates each correct response. So, if the number of error / contradictions equals or exceeds the number of marks available for the question, no marks can be awarded.

However, responses considered to be neutral (often prefaced by 'Ignore' in the mark scheme) are not penalised.

For example, in a question requiring 2 answers for 2 marks:

Correct answers	Incorrect answers (i.e. incorrect rather than neutral)	Mark (2)	Comment
1	0	1	
1	1	1	They have not exceeded the maximum number of responses so there is no penalty.
1	2	0	They have exceeded the maximum number of responses so the extra incorrect response cancels the correct one.
2	0	2	
2	1	1	
2	2	0	
3	0	2	The maximum mark is 2
3	1	1	The incorrect response cancels out one of the two correct responses that gained credit.
3	2	0	Two incorrect responses cancel out the two marks gained.
3	3	0	

3.2 Marking procedure for calculations

Full marks should be awarded for a correct numerical answer, without any working shown, unless the question states 'Show your working' or 'justify your answer'. In this case, the mark scheme will clearly indicate what is required to gain full credit.

If an answer to a calculation is incorrect and working is shown, process mark(s) can usually be gained by correct substitution / working and this is shown in the 'Comments' column or by each stage of a longer calculation.

3.3 Errors carried forward, consequential marking and arithmetic errors

Allowances for errors carried forward are most likely to be restricted to calculation questions and should be shown by the abbreviation ECF or consequential in the marking scheme.

An arithmetic error should be penalised for one mark only unless otherwise amplified in the marking scheme. Arithmetic errors may arise from a slip in a calculation or from an incorrect transfer of a numerical value from data given in a question.

3.4 Equations

In questions requiring students to write equations, state symbols are generally ignored unless otherwise stated in the 'Comments' column.

Examiners should also credit correct equations using multiples and fractions unless otherwise stated in the 'Comments' column.

3.5 Oxidation states

In general, the sign for an oxidation state will be assumed to be positive unless specifically shown to be negative.

3.6 Interpretation of 'it'

Answers using the word 'it' should be given credit only if it is clear that the 'it' refers to the correct subject.

3.7 Phonetic spelling

The phonetic spelling of correct scientific terminology should be credited **unless** there is a possible confusion with another technical term or if the question requires correct IUPAC nomenclature.

3.8 Brackets

(.....) are used to indicate information which is not essential for the mark to be awarded but is included to help the examiner identify the sense of the answer required.

3.9 Ignore / Insufficient / Do not allow

Ignore or insufficient is used when the information given is irrelevant to the question or not enough to gain the marking point. Any further correct amplification could gain the marking point.

Do **not** allow means that this is a wrong answer which, even if the correct answer is given, will still mean that the mark is not awarded.

3.10 Marking crossed out work

Crossed out work that **has not been** replaced should be marked as if it were not crossed out, if possible. Where crossed out work **has been** replaced, the replacement work and not the crossed out work should be marked.

3.11 Reagents

The command word “Identify”, allows the student to choose to use **either** the name or the formula of a reagent in their answer. In some circumstances, the list principle may apply when both the name and the formula are used. Specific details will be given in mark schemes.

The guiding principle is that a reagent is a chemical which can be taken out of a bottle or container. Failure to identify complete reagents **will be penalised**, but follow-on marks (e.g. for a subsequent equation or observation) can be scored from an incorrect attempt (possibly an incomplete reagent) at the correct reagent. Specific details will be given in mark schemes.

For example, **no credit** would be given for

- the cyanide ion or CN^- when the reagent should be potassium cyanide or KCN;
- the hydroxide ion or OH^- when the reagent should be sodium hydroxide or NaOH;
- the $\text{Ag}(\text{NH}_3)_2^+$ ion when the reagent should be Tollens' reagent (or ammoniacal silver nitrate). In this example, no credit is given for the ion, but credit could be given for a correct observation following on from the use of the ion. Specific details will be given in mark schemes.

In the event that a student provides, for example, **both** KCN and cyanide ion, it would be usual to ignore the reference to the cyanide ion (because this is not contradictory) and credit the KCN. Specific details will be given in mark schemes.


3.12 Organic structures

Where students are asked to draw organic structures, unless a specific type is required in the question and stated in the mark scheme, these may be given as displayed, structural or skeletal formulas or a combination of all three as long as the result is unambiguous.


In general

- Displayed formulae must show all of the bonds and all of the atoms in the molecule, but need not show correct bond angles.
- Skeletal formulae must show carbon atoms by an angle or suitable intersection in the skeleton chain. Functional groups must be shown and it is essential that all atoms other than C atoms are shown in these (except H atoms in the functional groups of aldehydes, secondary amines and N-substituted amides which do not need to be shown).
- Structures must not be ambiguous, e.g. 1-bromopropane should be shown as $\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}$ and not as the molecular formula $\text{C}_3\text{H}_7\text{Br}$ which could also represent the isomeric 2-bromopropane.
- Bonds should be drawn correctly between the relevant atoms. This principle applies in all cases where the attached functional group contains a carbon atom, e.g. nitrile, carboxylic acid, aldehyde and acid chloride. The carbon-carbon bond should be clearly shown. Wrongly bonded atoms will be penalised **on every occasion**. (see the examples below)
- The same principle should also be applied to the structure of alcohols. For example, if students show the alcohol functional group as $\text{C} - \text{HO}$, they should be penalised **on every occasion**.
- Latitude should be given to the representation of $\text{C} - \text{C}$ bonds in alkyl groups, given that CH_3- is considered to be interchangeable with $\text{H}_3\text{C}-$ even though the latter would be preferred.
- Similar latitude should be given to the representation of amines where $\text{NH}_2 - \text{C}$ will be allowed, although $\text{H}_2\text{N} - \text{C}$ would be preferred.
- Poor presentation of vertical $\text{C} - \text{CH}_3$ bonds or vertical $\text{C} - \text{NH}_2$ bonds should **not** be penalised. For other functional groups, such as $-\text{OH}$ and $-\text{CN}$, the limit of tolerance is the half-way position between the vertical bond and the relevant atoms in the attached group.


By way of illustration, the following would apply.




allowed




allowed




not allowed




not allowed




not allowed



allowed




allowed




allowed




allowed




not allowed




not allowed




not allowed




not allowed




not allowed




not allowed




not allowed




not allowed



not allowed



not allowed



not allowed

- Representation of CH₂ by C–H₂ will be penalised
- Some examples are given here of **structures** for specific compounds that should **not** gain credit (but, exceptions may be made in the context of balancing equations)

CH₃COH for ethanal

CH₃CH₂HO for ethanol

OHCH₂CH₃ for ethanol

C₂H₆O for ethanol

CH₂CH₂ for ethene

CH₂.CH₂ for ethene

CH₂:CH₂ for ethane

- Each of the following **should gain credit** as alternatives to correct representations of the structures.

CH₂ = CH₂ for ethene, H₂C=CH₂

CH₃CHOHCH₃ for propan-2-ol, CH₃CH(OH)CH₃

- In most cases, the use of “sticks” to represent C – H bonds in a structure should **not** be penalised. The exceptions to this when “sticks” will be penalised include
 - structures in mechanisms where the C – H bond is essential (e.g. elimination reactions in halogenoalkanes and alcohols)
 - when a displayed formula is required
 - when a skeletal structure is required or has been drawn by the candidate

3.13 Organic names


As a general principle, non-IUPAC names or incorrect spelling or incomplete names should **not** gain credit. Some illustrations are given here.

but-2-ol	should be butan-2-ol
2-hydroxybutane	should be butan-2-ol
butane-2-ol	should be butan-2-ol
2-butanol	should be butan-2-ol
ethan-1,2-diol	should be ethane-1,2-diol
2-methpropan-2-ol	should be 2-methylpropan-2-ol
2-methylbutan-3-ol	should be 3-methylbutan-2-ol
3-methylpentan	should be 3-methylpentane
3-mythylpentane	should be 3-methylpentane
3-methypentane	should be 3-methylpentane
propanitrile	should be propanenitrile
aminethane	should be ethylamine (although aminoethane can gain credit)
2-methyl-3-bromobutane	should be 2-bromo-3-methylbutane
3-bromo-2-methylbutane	should be 2-bromo-3-methylbutane
3-methyl-2-bromobutane	should be 2-bromo-3-methylbutane
2-methylbut-3-ene	should be 3-methylbut-1-ene
difluorodichloromethane	should be dichlorodifluoromethane


3.14 Organic reaction mechanisms

Curly arrows should originate either from a lone pair of electrons or from a bond.

The following representations should not gain credit **and will be penalised each time** within a clip.



For example, the following would score zero marks



When the curly arrow is showing the formation of a bond to an atom, the arrow can go directly to the relevant atom, alongside the relevant atom or **more than half-way** towards the relevant atom.

In free-radical substitution

- the absence of a radical dot should be penalised **once only** within a clip.
- the use of half-headed arrows is not required, but the use of double-headed arrows or the incorrect use of half-headed arrows in free-radical mechanisms should be penalised **once only** within a clip

The correct use of skeletal formulae in mechanisms is acceptable, but where a C-H bond breaks both the bond and the H must be drawn to gain credit.

Question	Marking guidance	Mark	Comments
01.1	$\text{C}_{15}\text{H}_{32} + 15\frac{1}{2} \text{O}_2 \longrightarrow 15\text{CO} + 16\text{H}_2\text{O}$	1	Allow multiples Ignore state symbols
01.2	Nitrogen and oxygen (from air) <u>react</u> At high temperatures	1 1	Allow correct equation Allow temperatures about 1000 °C / spark Not just heat/hot/heat from engine M2 dependent on M1
01.3	$\text{NO} + \text{CO} \longrightarrow \frac{1}{2} \text{N}_2 + \text{CO}_2$	1	Allow multiples Ignore state symbols
01.4	CaO/lime/CaCO ₃ /limestone $\text{CaCO}_3 + \text{SO}_2 \longrightarrow \text{CaSO}_3 + \text{CO}_2$ OR $\text{CaO} + \text{SO}_2 \longrightarrow \text{CaSO}_3$	1 1	Allow M1 scored from equation even if incorrect. Equation scores 2 marks Or via two equations: $\text{CaCO}_3 \longrightarrow \text{CaO} + \text{CO}_2$ $\text{CaO} + \text{SO}_2 \longrightarrow \text{CaSO}_3$ Allow $\text{SO}_2 + \text{CaO} + \frac{1}{2} \text{O}_2 \longrightarrow \text{CaSO}_4$ $\text{SO}_2 + \text{CaO} + 2\text{H}_2\text{O} + \frac{1}{2} \text{O}_2 \longrightarrow \text{CaSO}_4 + 2\text{H}_2\text{O}$ $\text{SO}_2 + \text{CaCO}_3 + \frac{1}{2} \text{O}_2 \longrightarrow \text{CaSO}_4 + \text{CO}_2$

01.5	High temperature / 700–800 K / 450 °C Zeolite catalyst / aluminosilicate	1	Ignore pressure
		1	Apply list principle “Catalyst” by itself is insufficient
01.6	$\text{C}_{15}\text{H}_{32} \longrightarrow \text{C}_6\text{H}_{14} + \text{C}_3\text{H}_6 + \text{C}_6\text{H}_{12}$	1	Allow multiples Ignore state symbols

Question	Marking guidance	Mark	Comments
02.1	$n(\text{Mg}) = \text{mass} \div A_r = 0.105 \div 24.3 = 0.00432$	1	Must be at least 2 sf
	$n(\text{HCl}) = n(\text{Mg}) \times 2 (= 0.00864)$	1	If incorrect ratio – lose M2 and M3
	$\text{vol} = 1000n/c = 1000 \times 0.00864 / 0.500 = 17.3 \text{ (cm}^3\text{)}$	1	Allow 17.28 (cm ³)
02.2	Tangent drawn at t=50 s Figure 2 Gradient of tangent calculated (must be at least 10 small squares in both x and y axes) – 0.65	1 1	Mark independently Allow range 0.61 – 0.69 Graph to be added at standardising


02.3	(Some) gas / hydrogen is lost before the syringe is connected to the flask	1	Penalise incorrect gas Not gas loss Allow “gas leaks”
02.4	Time taken would decrease as more particles in a given volume More collisions in a given time / more frequent collisions	1 1	If time increases CE=0

Question	Marking guidance	Mark	Comments
03.1	M1: The enthalpy change when 1 mol of a compound/substance is burned completely in oxygen / is burned in excess oxygen	1	
	M2: All substances in standard states	1	
03.2	$\Delta H = \Sigma \Delta H(\text{bonds broken}) - \Sigma \Delta H(\text{bonds formed})$	1	M2 for expression or substituted numbers
	$\Delta H =$	1	
	$7(\text{C-H}) + 2(\text{C-C}) + \text{C-O} + (\text{O-H}) + \overset{\mathbf{1}}{4\frac{1}{2}} (\text{O=O}) - [6(\text{C=O}) + 8(\text{O-H})]$ $-1886 =$ $7(412) + 2(348) + (\text{C-O}) + (463) + \overset{\mathbf{1}}{4\frac{1}{2}}(496) - [6(805) + 8(463)]$ $(\text{C-O}) = 373 \text{ (kJ mol}^{-1}\text{)}$	1	



03.3	$\Delta T = 18.0$ and mass of fuel = 0.566 g $q = 100 \times 4.18 \times 18 = 7524 \text{ J}$ $n = 0.566 \div 60.0 = 0.00943$ $\Delta H = -q \div n = -7.524 \div 0.00943 = -798 \text{ (kJ mol}^{-1}\text{)}$	1 1 1 1	M1 is for calculation of ΔT and mass of fuel M2 is for answer (q) M3 is for moles of fuel (3 sf or more) M4 is for a negative answer in kJ mol^{-1} Allow $-797.6 \text{ (kJ mol}^{-1}\text{)}$
03.4	Use copper can / use aluminium can (instead of glass beaker)	1	Allow add lid/add insulation on sides of container/ add draft shield/bomb calorimeter
03.5	$(2 \times 0.5) \div 16 \times 100 = 6.25(\%)$ or $6.3(\%)$	1	



Question	Marking guidance	Mark	Comments
04.1	Ag / silver	1	
04.2	C–O–C bond angle much smaller than 109.5°	1	Allow “bond angle only 60° ”
	Bonding pairs repel each other	1	Allow “Ring strain” / lower C-O bond enthalpy
04.3		4	

04.4					
	C	O	H		
	53.3/12.0	35.6/16.0	11.1/1.0	1	M1: divide by A_r
	4.44	2.23	11.1	1	M2: ratio 2:1:5
	2	1	5		
	Empirical formula = C_2OH_5				
	90/45 = 2, so Molecular formula = $C_4O_2H_{10}$			1	M3: molecular formula

Question	Marking guidance	Mark	Comments
05.1	2-methylbutanenitrile	1	
05.2	Nucleophilic substitution  (1) arrow + lp (1) arrow	1 1 1	M1 loss of Br- M2 attack of nucleophile on <u>correct</u> C+ Allow also Sn1 mechanism Allow structural formula
05.3	Base / proton acceptor	1	
	Elimination	1	Penalise “nucleophilic”
05.4	Slower (with 2-chlorobutane)	1	Allow converse. If M1 incorrect CE=0
	C-Cl bond stronger (than C-I)	1	C-I bond is weaker (than C-Cl)

05.5	D	1	
	C	1	
	B	1	

Question	Marking guidance	Mark	Comments
06.1	 M1: Arrow + lp M2: structure of ion M3 arrow M4 arrow	4	Allow E1: M2 and M3 separately via carbocation Allow M1 curly arrow to H of H ₂ SO ₄
06.2	Catalyst	1	Allow “to protonate the alcohol” Allow dehydrating agent
06.3		1	Ignore brackets Ignore n
06.4	Polymer does not contain double bond / polymer has a non-polar chain / Double bond in monomers are areas of high electron density / attract electrophiles	1	

Question	Marking guidance	Mark	Comments
07.1	X :  Y: methylcyclobutane Z: 2-methylbut-1-ene	1 1 1	
07.2	Test: Bromine (water) Observation with Y: no (visible) change Observation with Z: turns colourless / orange to colourless	1 1 1	If test is incorrect – CE=0 Not no reaction Allow correct alternative tests eg Test: Acidified KMnO_4 Observations with B: no (visible) change Observations with C: turns colourless / purple to colourless
07.3	 Atomic number of C > atomic number of H / H has lowest atomic number / H is always lowest priority group <u>Highest priority</u> atoms are on opposite sides for E isomer	1 1 1	If wrong isomer drawn allow M2 only

07.4	Two identical groups bonded to one carbon in the double bond / two hydrogen atoms on one carbon in the double bond / each of the carbons in the double bond is not bonded to two different (groups of) atoms	1	Must refer to C in a double bond or C=C
------	--	---	---

Question	Marking Guidance	Mark	Comments
08.1	Electrophilic addition	1	
08.2	2-bromopropane is the major product	1	Allow structure for M1
	Alkyl groups are electron releasing	1	
	Most stable carbocation is the secondary carbocation / the carbocation with the most alkyl groups attached to the C+	1	Must refer to carbocation as an intermediate