



Pearson  
Edexcel

Mark Scheme (Results)

January 2024

Pearson Edexcel International Advanced Level  
In Further Pure Mathematics F2 (WFM02)  
Paper 01

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

January 2024

Question Paper Log Number P73487A

Publications Code WFM02\_01\_2401\_MS

All the material in this publication is copyright

© Pearson Education Ltd 2024

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

# PEARSON EDEXCEL IAL MATHEMATICS

## General Instructions for Marking

1. The total number of marks for the paper is 75
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M** marks: Method marks are awarded for 'knowing a method and completing an attempt to apply it', unless otherwise indicated.
  - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.
3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- ft – follow through
- cao – correct answer only
- cso - correct solution only. There must be no clear errors in this part of the question to obtain this mark
- isw – ignore subsequent working
- awrt – answers which round to
- SC: special case
- oe – or equivalent
- dM – dependent method mark
- dp decimal places
- sf significant figures
- \* The answer is given on the paper – apply cso

4. All A marks are 'correct answer only' (cao), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.
5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
6. If a candidate makes more than one attempt at any question:
  - If all but one attempt is crossed out, mark the attempt which is NOT crossed out provided it is not cursory.
  - If either all attempts are crossed out or none are crossed out, score for their best attempt.
7. Ignore wrong working or incorrect statements following a correct answer unless the mark scheme indicates otherwise.
8. Mark question parts separately unless the scheme indicates otherwise.

**Usual rules for the method mark for solving a 3 term quadratic:**

**(Note: There may be schemes where the below does not apply)**

**If no method is shown then one root must be obtained that is consistent with their equation.**

**1. Factorisation**

$(x^2 + bx + c) = (x + p)(x + q)$ , where  $|pq| = |c|$ , leading to  $x = \dots$

$(ax^2 + bx + c) = (mx + p)(nx + q)$ , where  $|pq| = |c|$  and  $|mn| = |a|$ , leading to  $x = \dots$

**2. Formula**

Complete attempt to use the correct formula with values for a, b and c leading to  $x = \dots$  (may be unsimplified).

**3. Completing the square (where a = 1, otherwise must divide by a first - allow equivalent work if a is a square number)**

Solving  $x^2 + bx + c = 0$ :  $\left(x \pm \frac{b}{2}\right)^2 \pm q \pm c = 0$ ,  $q \neq 0$ , leading to  $x = \dots$

**January 2024**  
**WFM02 Further Pure Mathematics F2 Mark Scheme**

Question Number	Scheme	Notes	Marks	
<b>1</b>	$\frac{1}{x+2} > 2x+3$			
	<p style="text-align: center;">Examples:</p> $\frac{1-(x+2)(2x+3)}{x+2} > 0 \Rightarrow 2x^2 + 7x + 5 = 0$ $x+2 > (2x+3)(x+2)^2$ $\Rightarrow (x+2)(2x^2 + 7x + 5) = 0 \text{ or } 2x^3 + 11x^2 + 19x + 10 = 0$ $\frac{1}{x+2} = 2x+3 \Rightarrow (2x+3)(x+2) - 1 = 2x^2 + 7x + 5 = 0$ <p>Uses algebra to obtain a 3TQ, <math>(x+2)</math> multiplied by a 3TQ or a 4TC. Allow slips and condone incorrect inequality signs but the first algebraic step should be otherwise appropriate so do not accept work with e.g., <math>(2x+3)(x+2) = 0</math>. The “= 0” can be implied by solutions. Graphical attempts require intersections to be found algebraically. Squaring first is acceptable so allow M1 for obtaining a 5TQ <math>(4x^4 + 28x^3 + 73x^2 + 84x + 35 = 0)</math></p>		<b>M1</b>	
	e.g., $(2x+5)(x+1) = 0 \Rightarrow$ $x = -\frac{5}{2}, -1$	Both $-1$ and $-\frac{5}{2}$ from appropriate work and no extra incorrect cvs. May only be seen in the solution set. Allow solving a 3TQ etc. by calculator.		<b>A1</b>
	$x = -2$	Identifies $-2$ as a critical value. May only be seen in solution set. This is the only mark available if there is no algebraic manipulation seen. Allow from any or no working e.g., from $(2x+3)(x+2) = 0$		<b>B1</b>
	$\Rightarrow x < -\frac{5}{2}, -2 < x < -1$ or e.g., $(-\infty, -2.5), (-2, -1)$			<b>M1 A1</b>
	<p><b>M1:</b> For the regions <math>x &lt; a, -2 &lt; x &lt; b</math> with real cvs <math>a &lt; -2</math> and <math>b &gt; -2</math> <b>but condone</b> <math>b &lt; x &lt; -2</math> as a notational slip for this mark.</p> <p>Condone any non-strict inequality signs and poor notation for this mark. Not dependent but must follow an attempt at algebraic manipulation.</p> <p><b>A1:</b> Correct solution set in any form. Do not isw if the correct inequalities are subsequently incorrectly amended. Allow all marks even if an incorrect inequality sign was seen earlier in the working.</p>			
	<p style="text-align: center;">Examples:</p> $-\frac{5}{2} > x \text{ or } -2 < x < -1 \text{ M1 A1} \quad x < -\frac{5}{2} \text{ and } -2 < x < -1 \text{ M1 A1}$ <p style="text-align: center;">(Accept any word between the two correct regions)</p> $x < -\frac{5}{2}, -1 < x < -2 \text{ M1 A0 (notational slip)}$ $\left(-\infty, -\frac{5}{2}\right) \cap (-2, -1) \text{ M1A0 (incorrect symbol – allow “and”)} \quad \left[-\infty, -\frac{5}{2}\right] \cup [-2, -1] \text{ M1A0}$ $x < -\frac{5}{2} \quad -2 < x \quad x < -1 \text{ M0 A0 (insufficient)}$			
			<b>(5)</b>	
			<b>Total 5</b>	

Question Number	Scheme	Notes	Marks
<b>2(a)</b>	<b>(i)</b> $z = 6 - 6\sqrt{3}i \Rightarrow  z  = \sqrt{6^2 + (6\sqrt{3})^2} = 12$	+12 only. Accept if just stated	<b>B1</b>
	<b>(ii)</b> e.g., $\arg z = -\arctan \frac{6\sqrt{3}}{6}$ Attempts an expression for a relevant angle. Look for $\pm \arctan \left( \pm \frac{6\sqrt{3}}{6} \right)$ or e.g., $\pm \tan^{-1} \left( \pm \frac{1}{\sqrt{3}} \right)$ If arctan is not seen allow e.g., $\tan \alpha = \frac{6\sqrt{3}}{6} \Rightarrow \alpha = \frac{\pi}{3}$ with $\alpha$ correct for their $\tan \alpha$ If using sin or cos the hypotenuse must be their 12		<b>M1</b>
	$\arg z$ or arg or argument (of $z$ ) $= -\frac{\pi}{3}$ * A correct proof with no incorrect work/statements. LHS required. <b>Allow "θ = "</b> if <b>consistent</b> , e.g., $\theta = -\frac{\pi}{3}$ cannot follow " $\tan \theta = +\sqrt{3}$ "		<b>A1*</b>
<b>(ii)</b> <b>Way 2</b>	$z = 12\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right) = 12\left(\cos\left(-\frac{\pi}{3}\right) + i\sin\left(-\frac{\pi}{3}\right)\right)$ or $12e^{-\frac{\pi}{3}i}$ or $\cos \theta = \frac{1}{2}$ or $\sin \theta = -\frac{\sqrt{3}}{2}$ [M1] $\Rightarrow \arg z = -\frac{\pi}{3}$ [A1*] M1: Factorises out 12 and writes in trig or exp form or identifies $\cos \theta = \frac{1}{2}$ and $\sin \theta = -\frac{\sqrt{3}}{2}$ A1: Acceptable statement with all work correct		
<b>(ii)</b> <b>Way 3</b>	$z = 12\left(\cos\left(-\frac{\pi}{3}\right) + i\sin\left(-\frac{\pi}{3}\right)\right)$ or $12e^{-\frac{\pi}{3}i}$ or $12\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right) = 6 - 6\sqrt{3}i$ [M1] $\Rightarrow \arg z = -\frac{\pi}{3}$ [A1*] M1: Assumes result, writes correctly for their 12 and attempts $a + ib$ form A1: Obtains $6 - 6\sqrt{3}i$ and makes acceptable statement with all work correct		
			<b>(3)</b>
<b>(b)</b>	$z = "12" \left( \cos\left(-\frac{\pi}{3}\right) + i\sin\left(-\frac{\pi}{3}\right) \right)$ or $"12" e^{-\frac{\pi}{3}i}$ [no missing "i" unless recovered] Correct trig or exp. form with their 12. Could be implied by their $z^4$ in trig or exp. form e.g., $(\text{"12" } e^{-\frac{\pi}{3}i})^4$ Allow equivalent values of $\theta$ e.g. $\frac{5\pi}{3}$ and use of e.g., $\sin\left(-\frac{\pi}{3}\right) = -\sin\left(\frac{\pi}{3}\right)$ . Condone poor bracketing. Allow this mark if $+2k\pi, -2k\pi, \pm 2k\pi$ appears with argument		<b>M1</b>
	$z^4 = 20736 \left( \cos\left(-\frac{4\pi}{3}\right) + i\sin\left(-\frac{4\pi}{3}\right) \right)$ or $20736 \left( \cos -\frac{4\pi}{3} + i\sin -\frac{4\pi}{3} \right)$ or $20736 e^{-\frac{4\pi}{3}i}$ Correct $z^4$ in any form. $12^4$ evaluated and arg. of $-\frac{4\pi}{3}$ (not just $4 \times -\frac{\pi}{3}$ ) or $\frac{2\pi}{3}$ only although may use e.g., $\sin\left(-\frac{4\pi}{3}\right) = -\sin\left(\frac{4\pi}{3}\right)$ . No "k"s. Condone an "unclosed" bracket. Only accept $-10368 + 10368\sqrt{3}i$ or $20736\left(-\frac{1}{2} + \frac{\sqrt{3}}{2}i\right)$ <b>provided evidence of de Moivre.</b>		<b>A1</b>
			<b>(2)</b>

Question Number	Scheme	Notes	Marks
2(c)	$w = z^{\frac{1}{2}} = (\pm)\sqrt[12]{12} \left( \cos\left(\frac{-\pi}{3}\right) + i \sin\left(\frac{-\pi}{3}\right) \right)$ <p style="text-align: center;">[no missing "i" unless recovered]</p> <p>Correct use of de Moivre's theorem with <math>-\frac{\pi}{3}</math> and their 12 to attempt one square root.</p> <p>Allow work with argument of <math>\frac{5\pi}{3}</math> for <math>-\frac{\pi}{3}</math> and use of e.g., <math>\sin\left(-\frac{\pi}{6}\right) = -\sin\left(\frac{\pi}{6}\right)</math>. Condone poor bracketing.</p> <p>M0 if <math>z^4</math> used for <math>z</math>. Allow this mark if <math>+2k\pi, -2k\pi, \pm 2k\pi</math> appears with argument</p>		<b>M1</b>
	$w = 3 - \sqrt{3}i, -3 + \sqrt{3}i \text{ oe}$ <p>A1ft: One correct exact root in <math>a + ib</math> or <math>c(a + ib)</math> form (<math>a, b, c</math> may be unsimplified but not numerical trig expressions) <b>ft their 12 only</b> i.e. <math>(\pm)\sqrt[12]{12} \left( \frac{\sqrt{3}}{2} - \frac{1}{2}i \right)</math></p> <p>A1: Both exact roots (no others) correct in <math>a + ib</math> form – <math>a</math> and <math>b</math> may be unsimplified (but not numerical trig expressions) e.g. accept</p> $a = (\pm) \sqrt{12} \frac{\sqrt{3}}{2}, (\pm) \frac{\sqrt{36}}{2} \quad b = (\mp) \frac{\sqrt{12}}{2}, (\mp) \frac{2\sqrt{3}}{2}$ <p>Accept <math>\pm(3 - \sqrt{3}i)</math> but just <math>\pm 3 - \sqrt{3}i</math> is A1 A0. Just <math>\pm\sqrt{3}(\sqrt{3} - i)</math> is A1 A0</p>		
	Note: $w^2 = r^2(\cos 2\theta + i \sin 2\theta) = z \Rightarrow r, \theta, w = \dots$ is an acceptable approach		<b>(3)</b>
<b>Alt</b>	$w^2 = z \Rightarrow (a + ib)^2 = a^2 - b^2 + 2abi = 6 - 6\sqrt{3}i \Rightarrow a^2 - b^2 = 6, 2ab = -6\sqrt{3}$ $b = -\frac{3\sqrt{3}}{a} \Rightarrow a^2 - \frac{27}{a^2} = 6 \Rightarrow a^4 - 6a^2 - 27 = (a^2 - 9)(a^2 + 3) = 0 \Rightarrow a^2 = 9, a = \pm 3, b = \mp\sqrt{3}$ <p><b>M1:</b> From a correct starting point, expands and equates real and imaginary parts to form two equations in <math>a</math> and <math>b</math> and obtains at least one value for both <math>a</math> and <math>b</math></p> $w = 3 - \sqrt{3}i, -3 + \sqrt{3}i$ <p><b>A1:</b> One correct exact root in <math>a + ib</math> or <math>c(a + ib)</math> form (<math>a, b, c</math> may be unsimplified)</p> <p><b>A1:</b> Both exact roots (no others) correct in <math>a + ib</math> form – <math>a</math> and <math>b</math> may be unsimplified</p>		
			<b>Total 8</b>



Question Number	Scheme	Notes	Marks
<b>3(a)</b>	$\frac{r}{\sqrt{r(r+1)} + \sqrt{r(r-1)}} \times \frac{\sqrt{r(r+1)} - \sqrt{r(r-1)}}{\sqrt{r(r+1)} - \sqrt{r(r-1)}}$	A correct multiplier to rationalise the denominator seen or implied by correct work	<b>M1</b>
	$= \frac{r(\sqrt{r(r+1)} - \sqrt{r(r-1)})}{r(r+1) - r(r-1)} = \frac{\sqrt{r(r+1)} - \sqrt{r(r-1)}}{2} \text{ or } A = \frac{1}{2}$ <p>Correct expression or correct value for A. Condone poor notation if intention clear. There must be (minimal) correct supporting working.</p>		<b>A1</b>
	<p>Alternative:</p> $A = \frac{r}{(\sqrt{r(r+1)} + \sqrt{r(r-1)})(\sqrt{r(r+1)} - \sqrt{r(r-1)})} = \frac{r}{r(r+1) - r(r-1)} \text{ or } \frac{r}{r^2 + r - r^2 + r} \text{ or } \frac{r}{2r} \Rightarrow A = \frac{1}{2}$ <p>M1: Correctly makes A the subject A1: Correct completion with one intermediate fraction</p>		
			<b>(2)</b>
<b>(b)</b>	$\sum_{r=1}^n \frac{r}{\sqrt{r(r+1)} + \sqrt{r(r-1)}} = \frac{1}{2} \left( \begin{array}{l} \sqrt{1 \times 2} - \sqrt{1 \times 0} (= \sqrt{2} (-0)) \\ + \sqrt{2 \times 3} - \sqrt{2 \times 1} (= \sqrt{6} - \sqrt{2}) + \dots \\ \dots + \sqrt{(n-1)(n-1+1)} - \sqrt{(n-1)(n-1-1)} (= \sqrt{n(n-1)} - \sqrt{(n-1)(n-2)}) \\ + \sqrt{n(n+1)} - \sqrt{n(n-1)} \end{array} \right)$ <p><b>M1:</b> Applies the method of differences for <math>r=1</math> and <math>r=n</math> in the given expression with or without their A and obtains one correct row of these 2.</p> <p><b>M1:</b> Applies the method of differences for <math>r=1</math>, <math>r=n</math> and <b>either</b> <math>r=2</math> or <math>r=n-1</math> in the given expression with/without their A and obtains 2 correct rows of these 4.</p> <p>When considering how many rows are correct, if A has been clearly applied to any term then assess all rows as if A has been applied throughout.</p> <p>Condone missing bracket if their A is applied to a row e.g., "<math>\frac{1}{2} \times \sqrt{6} - \sqrt{2}</math>" <u>if it is recovered</u> but e.g., <math>\frac{\sqrt{6}}{2} - \sqrt{2}</math> is an incorrect row. Ignore a row for <math>r=0</math>. Condone equivalent work with <math>r</math> or e.g., <math>k</math> used for <math>n</math>.</p> <p><b>Both marks can be implied by a correct final expression with or without their A provided there are at least any two correct rows of differences</b></p> <p>i.e., "<math>\frac{1}{2}(\sqrt{n(n+1)} - 0)</math>" or <math>\sqrt{n(n+1)} - 0</math></p> <p>Note: row 3 is "<math>\frac{1}{2}(\sqrt{12} \text{ (or } 2\sqrt{3}) - \sqrt{6})</math>", row 4 is "<math>\frac{1}{2}(\sqrt{20} \text{ (or } 2\sqrt{5}) - \sqrt{12} \text{ (or } 2\sqrt{3}))</math>"</p> <p>If <math>\frac{1}{2}</math> is fully applied the rows are:</p> $\frac{\sqrt{2}}{2}, \frac{\sqrt{6}}{2} - \frac{\sqrt{2}}{2}, \frac{\sqrt{12}}{2} \text{ (or } \sqrt{3}) - \frac{\sqrt{6}}{2}, \frac{\sqrt{20}}{2} \text{ (or } \sqrt{5}) - \frac{\sqrt{12}}{2} \text{ (or } \sqrt{3}), \dots$ $\dots, \frac{\sqrt{(n-2)(n-1)}}{2} - \frac{\sqrt{(n-2)(n-3)}}{2}, \frac{\sqrt{n(n-1)}}{2} - \frac{\sqrt{(n-1)(n-2)}}{2}, \frac{\sqrt{n(n+1)}}{2} - \frac{\sqrt{n(n-1)}}{2}$	<p>Correct expression in terms of <math>n</math>. No incorrect terms seen in differences work even if cancelled but condone the occasional poor bracket. There should be no "0" so e.g.,</p> $\frac{1}{2}(\sqrt{n(n+1)} - 0) \text{ is A0}$ <p>Does not require marks in (a)</p>	<b>M1</b> <b>M1</b>
	$= \frac{1}{2} \sqrt{n(n+1)} \text{ oe e.g., } \frac{\sqrt{n^2 + n}}{2}$		<b>A1</b>
			<b>(3)</b>

Question Number	Scheme	Notes	Marks
<b>3(c)</b>	$\sum r = \frac{1}{2}n(n+1)$ e.g., sight of $k \times \dots = \sqrt{\frac{1}{2}n(n+1)}$	States or uses the correct summation formula for integers	<b>M1</b>
	$\frac{k}{2}\sqrt{n(n+1)} = \sqrt{\frac{1}{2}n(n+1)} \Rightarrow \frac{k}{2} = \sqrt{\frac{1}{2}} \Rightarrow k = \sqrt{2}$	$\sqrt{2}$ only (Not $\pm$ ). $k = \sqrt{2}$ must not come from a clearly incorrect equation.	<b>A1</b>
			<b>(2)</b>
			<b>Total 7</b>

Question Number	Scheme	Notes	Marks
<b>4(a)</b>	$y = \tan\left(\frac{3x}{2}\right) \Rightarrow y' = \frac{3}{2} \sec^2\left(\frac{3x}{2}\right)$	Any correct first derivative. Not implied by $y'\left(\frac{\pi}{6}\right) = 3$	<b>B1</b>
	$\Rightarrow y'' = 2 \times \frac{3}{2} \sec\left(\frac{3x}{2}\right) \times \sec\left(\frac{3x}{2}\right) \tan\left(\frac{3x}{2}\right) \times \frac{3}{2}$ $\left[ = \frac{9}{2} \sec^2\left(\frac{3x}{2}\right) \tan\left(\frac{3x}{2}\right) \right]$	Attempts the second derivative achieving $k \sec^2\left(\frac{3x}{2}\right) \tan\left(\frac{3x}{2}\right)$ or unsimplified equivalent. Not implied by $y''\left(\frac{\pi}{6}\right) = 9$	<b>M1</b>
	$\Rightarrow y''' = \frac{9}{2} \sec^2\left(\frac{3x}{2}\right) \sec^2\left(\frac{3x}{2}\right) \times \frac{3}{2} + \frac{9}{2} \tan\left(\frac{3x}{2}\right) \times 2 \times \frac{3}{2} \sec^2\left(\frac{3x}{2}\right) \tan\left(\frac{3x}{2}\right)$ $\left[ = \frac{27}{4} \sec^4\left(\frac{3x}{2}\right) + \frac{27}{2} \sec^2\left(\frac{3x}{2}\right) \tan^2\left(\frac{3x}{2}\right) \right]$	<b>dM1:</b> Attempts third derivative using the product rule, achieving $P \sec^4\left(\frac{3x}{2}\right) + Q \sec^2\left(\frac{3x}{2}\right) \tan^2\left(\frac{3x}{2}\right)$ or unsimplified equivalent. <b>Requires previous M mark.</b> <b>A1:</b> Correct differentiation. Accept unsimplified. Not implied by $y'''\left(\frac{\pi}{6}\right) = 54$	<b>dM1</b> <b>A1</b>
	If $\sec^2\left(\frac{3x}{2}\right) = \tan^2\left(\frac{3x}{2}\right) + 1$ is used the identity must be used correctly and to score M marks expressions of consistent form should be achieved. Note that replacing $\sec^2\left(\frac{3x}{2}\right)$ in $y'' \Rightarrow y''' = \frac{27}{4} \sec^2\left(\frac{3x}{2}\right) + \frac{81}{4} \sec^2\left(\frac{3x}{2}\right) \tan^2\left(\frac{3x}{2}\right)$		
	$y\left(\frac{\pi}{6}\right) = 1, y'\left(\frac{\pi}{6}\right) = 3, y''\left(\frac{\pi}{6}\right) = 9, y'''\left(\frac{\pi}{6}\right) = 54$ Attempts values (but allow numerical trig expressions) for y and their 3 derivatives at $\frac{\pi}{6}$ - accept stated values or insertion into a series of the correct form		
	$(y =) 1 + 3\left(x - \frac{\pi}{6}\right) + \frac{9}{2!}\left(x - \frac{\pi}{6}\right)^2 + \frac{54}{3!}\left(x - \frac{\pi}{6}\right)^3 + \dots$ Applies Taylor's correctly about $\frac{\pi}{6}$ with their values/numerical trig expressions. If values are not seen separately the work should imply a correct formula but allow a recognisable attempt at the series following the correct general formula stated. <b>Requires previous M mark.</b>		
	$(y =) 1 + 3\left(x - \frac{\pi}{6}\right) + \frac{9}{2}\left(x - \frac{\pi}{6}\right)^2 + 9\left(x - \frac{\pi}{6}\right)^3 + \dots$	Correct expression with coeffs. in simplest form. "y = ..." not required. <b>Requires all previous marks.</b> Score A0 if <b>clear evidence</b> of <u>use</u> of any wrong derivative expression.	<b>A1</b>
If e.g. $y'''\left(\frac{\pi}{6}\right)$ is found by calculator but $y'(x)$ and $y''(x)$ were seen award 1100110 max			<b>(7)</b>
Note: With responses that work in sin and cos throughout, to score M marks there must be no loss of form when differentiating (sign and coefficient errors only, also allowing sign errors with product/quotient formulae). Any use of identities must be correct. E.g: $y = \tan\left(\frac{3x}{2}\right) = \frac{\sin\left(\frac{3x}{2}\right)}{\cos\left(\frac{3x}{2}\right)} \Rightarrow y' = \frac{\frac{3}{2} \cos^2\left(\frac{3x}{2}\right) + \frac{3}{2} \sin^2\left(\frac{3x}{2}\right)}{\cos^2\left(\frac{3x}{2}\right)}$ $y'' = \frac{\frac{9}{2} \cos^3\left(\frac{3x}{2}\right) \sin\left(\frac{3x}{2}\right) + \frac{9}{2} \cos\left(\frac{3x}{2}\right) \sin^3\left(\frac{3x}{2}\right)}{\cos^4\left(\frac{3x}{2}\right)} \text{ or } \frac{\frac{9}{2} \cos\left(\frac{3x}{2}\right) \sin\left(\frac{3x}{2}\right)}{\cos^4\left(\frac{3x}{2}\right)} \text{ or } \frac{9 \sin\left(\frac{3x}{2}\right)}{2 \cos^3\left(\frac{3x}{2}\right)}$ $y''' = \frac{\frac{27}{4} \cos^8\left(\frac{3x}{2}\right) + 27 \cos^6\left(\frac{3x}{2}\right) \sin^2\left(\frac{3x}{2}\right) + \frac{81}{4} \cos^4\left(\frac{3x}{2}\right) \sin^4\left(\frac{3x}{2}\right)}{\cos^8\left(\frac{3x}{2}\right)} = \frac{27}{4} + 27 \tan^2\left(\frac{3x}{2}\right) + \frac{81}{4} \tan^4\left(\frac{3x}{2}\right)$			

Question Number	Scheme	Notes	Marks
<b>4(b)</b>	$\left\{ y\left(\frac{\pi}{4}\right) = \right\} 1 + 3\left(\frac{\pi}{4} - \frac{\pi}{6}\right) + \frac{9}{2}\left(\frac{\pi}{4} - \frac{\pi}{6}\right)^2 + 9\left(\frac{\pi}{4} - \frac{\pi}{6}\right)^3$ $\text{or } 1 + 3\left(\frac{\pi}{12}\right) + \frac{9}{2}\left(\frac{\pi}{12}\right)^2 + 9\left(\frac{\pi}{12}\right)^3$ <p>Substitutes <math>\frac{\pi}{4}</math> into their expression for <math>y</math> of the correct form with at least the first three terms (series about <math>\frac{\pi}{6}</math>). Must have values (not unevaluated trig expressions).  If only a decimal value is given then it must be the correct awrt 2.26 to score M1 (2.255314325).  If there is no working they must obtain an expression with at least <math>a + b\pi + c\pi^2</math> and correct exact ft <math>a</math>, <math>b</math> and <math>c</math> for their series or <math>1 + \frac{\pi}{4} + c\pi^2</math> with correct exact ft <math>c</math></p>		<b>M1</b>
	$= 1 + \frac{\pi}{4} + \frac{\pi^2}{32} + \frac{\pi^3}{192} \text{ or } 1 + \frac{1}{4}\pi + \frac{1}{32}\pi^2 + \frac{1}{192}\pi^3$	Correct answer or values for $A$ (32) and $B$ (192). Can be awarded if full marks were not scored in (a).	<b>A1</b>
		<b>(2)</b>	
		<b>Total 9</b>	

Question Number	Scheme	Notes	Marks
5	$r^2 = 100\cos^2 \theta + 20\cos \theta \tan \theta + \tan^2 \theta$	Any correct expression for $r^2$	<b>B1</b>
	$\left\{ \frac{1}{2} \right\} \int_0^{\frac{\pi}{3}} r^2 d\theta = \left\{ \frac{1}{2} \right\} \int_0^{\frac{\pi}{3}} (100\cos^2 \theta + 20\sin \theta + \tan^2 \theta) \{d\theta\}$	Attempts formula for the area with their $r^2$ which may not be expanded Condone missing $\frac{1}{2}$ and limits not required	<b>M1</b>
	$= \frac{1}{2} \int_0^{\frac{\pi}{3}} (50(1 + \cos 2\theta) + 20\sin \theta + \sec^2 \theta - 1) \{d\theta\}$ <b>M1:</b> Uses $\cos^2 \theta = \pm \frac{1}{2} \pm \frac{1}{2} \cos 2\theta$ <b>or</b> $\tan^2 \theta = \pm \sec^2 \theta \pm 1$ in their $r^2$ <b>M1:</b> Uses both $\cos^2 \theta = \pm \frac{1}{2} \pm \frac{1}{2} \cos 2\theta$ <b>and</b> $\tan^2 \theta = \pm \sec^2 \theta \pm 1$ in their $r^2$ Both M marks can be scored without the integral and the $\frac{1}{2}$ . Condone mixed variables. <b>A1:</b> Correct integral following $\cos^2 \theta = \frac{1}{2} + \frac{1}{2} \cos 2\theta$ and $\tan^2 \theta = \sec^2 \theta - 1$ . The $\cos \theta \tan \theta$ must be written as $\sin \theta$ (implied if appropriately integrated later). The $\frac{1}{2}$ is required (it may be seen later) but limits/ $d\theta$ are not needed. Allow mixed variables if subsequent work recovers this.	<b>M1</b> <b>M1</b> <b>A1</b>	
	$= \frac{1}{2} \left[ 49\theta + 25\sin 2\theta - 20\cos \theta + \tan \theta \right]_0^{\frac{\pi}{3}} \text{ or } \left[ \frac{49}{2}\theta + \frac{25}{2}\sin 2\theta - 10\cos \theta + \frac{1}{2}\tan \theta \right]_0^{\frac{\pi}{3}}$ <b>M1:</b> Achieves <b>three</b> of the following four integrated forms: $k \rightarrow k\theta$ (at least once), $\cos 2\theta \rightarrow \dots \sin 2\theta$ , $\sin \theta \rightarrow \dots \cos \theta$ , $\sec^2 \theta \rightarrow \dots \tan \theta$ . Ignore other terms if 3 of the above are satisfied. No $\frac{1}{2}$ or limits required. Condone mixed variables. <b>A1:</b> Correct integration including the $\frac{1}{2}$ (may be seen later). Limits not required. May be unsimplified e.g., $49\theta$ seen as $50\theta - \theta$ . Allow mixed variables if subsequent work recovers this.	<b>M1</b> <b>A1</b>	
	$= \frac{1}{2} \left( \frac{49\pi}{3} + 25\sin \frac{2\pi}{3} - 20\cos \frac{\pi}{3} + \tan \frac{\pi}{3} - (0 + 0 - 20 + 0) \right)$ $\left\{ = \frac{1}{2} \left( \frac{49\pi}{3} + \frac{25\sqrt{3}}{2} - 10 + \sqrt{3} + 20 \right) \text{ or } \frac{49\pi}{6} + \frac{25\sqrt{3}}{4} - 5 + \frac{\sqrt{3}}{2} + 10 \right\}$ Applies the correct limits to an expression of the form $p\theta + q \sin 2\theta + r \cos \theta + s \tan \theta$ ( $p, q, r, s \neq 0$ ) Allow slips but there must be a clear attempt to substitute, and they must only subtract the value of their $r$ , e.g. if $r = -20$ work must have or imply $\dots - (-20)$ or $+20$ . Allow mixed variables if the substitution recovers this.	<b>M1</b>	
$= \frac{1}{12} (98\pi + 81\sqrt{3} + 60)$	Correct answer or values for $a, b$ & $c$	<b>A1</b>	
Note that there are other viable routes through the integration e.g., use of integration by parts			<b>(9)</b>
			<b>Total 9</b>


Question Number	Scheme	Notes	Marks
6	$\frac{d^2x}{dt^2} + 6\frac{dx}{dt} + 13x = 8e^{-3t}$	$t \dots 0$	
(a)	$m^2 + 6m + 13 = 0 \Rightarrow m = \frac{-6 \pm \sqrt{36 - 52}}{2}$ $\{ = -3 \pm 2i \}$	Forms correct auxiliary equation and obtains a correct numerical expression for at least one root by formula or uses CTS (apply usual CTS rule below). One correct root if no working	<b>M1</b>
	CTS rule: $m^2 + 6m + 13 = 0 \Rightarrow \left(m \pm \frac{6}{2}\right)^2 \pm q \pm 13 = 0, q \neq 0 \Rightarrow m = \dots$		
	CF examples: $(x =) e^{-3t} (A \cos 2t + B \sin 2t)$ or $(x =) Ae^{-3t} \cos(-2t) + Be^{-3t} \sin(-2t)$ or $(x =) Pe^{(-3+2i)t} + Qe^{(-3-2i)t}$ or $(x =) e^{-3t} (Pe^{2it} + Qe^{-2it})$	Correct complementary function in any form, allow if the "x =" is missing or wrong and accept for this mark if the CF is given fully in terms of x instead of t.	<b>A1</b>
	PI: $\{x =\} \lambda e^{-3t}$	Correct form for the particular integral selected. Must include $\lambda e^{-3t}$ but accept with any extra terms that correctly disappear when coefficients found. Accept "PI=". If $\lambda e^{pt}$ is used $p = -3$ must be seen later.	<b>B1</b>
	$\frac{dx}{dt} = -3\lambda e^{-3t}; \frac{d^2x}{dt^2} = 9\lambda e^{-3t}$ $\Rightarrow 9\lambda e^{-3t} + 6(-3\lambda e^{-3t}) + 13\lambda e^{-3t} = 8e^{-3t}$	Differentiates a PI of <b>any</b> form twice ( <b>provided</b> it has at least one constant and is a function of t) <b>and</b> substitutes into the equation. Allow only sign/coefficient errors only in the differentiation. Their PI must lead to non-zero derivatives.	<b>M1</b>
	$\Rightarrow 9\lambda - 18\lambda + 13\lambda = 8 \Rightarrow \lambda = \dots (2)$	Proceeds to find the value of the constant <b>following use of a PI of the correct form</b> . Any unnecessary extra terms in the PI must be found to be zero	<b>dM1</b>
	$x = "e^{-3t} (A \cos 2t + B \sin 2t)" + 2e^{-3t}$	Correct general solution ft on <b>their CF only</b> – any CF <b>provided</b> it has at least one constant and is in terms of t. <b>Must have x = ...</b> Do not allow if their CF is miscopied or mathematically changed	<b>A1ft</b>
Work with a PI of the form $\lambda te^{-3t}$ is BOM1dM0A0 max even if $2e^{-3t}$ is obtained. Only condone incorrect variables if they are recovered but refer to the note for the first A1.			<b>(6)</b>

Question Number	Scheme	Notes	Marks
6(b)	$x = \frac{1}{2} \text{ at } t = 0$ $\Rightarrow \frac{1}{2} = A + 2 \quad \left( \Rightarrow A = -\frac{3}{2} \right)$	Uses the initial condition for $x$ in their GS to find a linear equation in one or two constants. Allow for GS = CF or CF + PI and the constant may come from the +PI	M1
	$x = e^{-3t} (A \cos 2t + B \sin 2t) + 2e^{-3t}$ $\frac{dx}{dt} = e^{-3t} (-2A \sin 2t + 2B \cos 2t) - 3e^{-3t} (A \cos 2t + B \sin 2t) - 6e^{-3t}$ <p>Uses the <b>product rule</b> to differentiate their real GS obtaining an expression in terms of <math>t</math> of the correct form for their GS (sign and coefficient errors only – so do not allow e.g., <math>\dots e^{pt} \rightarrow \dots e^{qt}</math>). Allow for GS = CF or CF + PI and does not have to include constants.</p> <p>If they work with a complex function e.g., <math>x = Pe^{(-3+2i)t} + Qe^{(-3-2i)t} + 2e^{-3t}</math> progress is unlikely.</p> <p>This mark is not scored for work in (c)</p>		M1
	$t = 0, \frac{dx}{dt} = \frac{1}{2} \Rightarrow \frac{1}{2} = 2B - 3A - 6 \Rightarrow B = \dots (=1)$ <p>Uses both initial conditions to find values for the 2 constants (no others) in their <b>GS = (CF with 2 constants) + PI(no constants)</b>. One constant must be found to be non-zero.</p> <p><b>Requires both previous M marks.</b></p>		ddM1
	<p>Examples:</p> $x = e^{-3t} \left( -\frac{3}{2} \cos 2t + \sin 2t \right) + 2e^{-3t}$ $\text{or } x = e^{-3t} \left( -\frac{3}{2} \cos 2t + \sin 2t + 2 \right)$ $\text{or } x = 2e^{-3t} - \frac{3}{2} e^{-3t} \cos 2t + e^{-3t} \sin 2t$	Correct particular solution in any form in terms of $t$ . <b>Must be <math>x = \dots</math> unless this was the only reason for final A0 in part (a) due to omission or e.g, “<math>y = \dots</math>” was used</b>	A1
			(4)
(c)	$\frac{dx}{dt} = e^{-3t} (3 \sin 2t + 2 \cos 2t) - 3e^{-3t} \left( -\frac{3}{2} \cos 2t + \sin 2t \right) - 6e^{-3t} = 0$ <p>Sets an expression for <math>\frac{dx}{dt} = 0</math>. Accept with any unfound constants provided <math>\frac{dx}{dt} = f(t)</math></p>		M1
	$(3 \sin 2t + 2 \cos 2t) - 3 \left( -\frac{3}{2} \cos 2t + \sin 2t \right) - 6 = 0$ <p>Achieves an equation of the form <math>a \sin bt + c \cos bt + d = 0</math> or equivalent with terms uncollected. One of <math>a</math> and <math>c</math> non-zero and <math>b</math> and <math>d</math> non-zero.</p> <p><b>Must follow a GS = CF + PI where two constants were found for the CF and one for the PI. Requires previous M mark.</b></p>		dM1
	$\cos 2t = \frac{12}{13} \Rightarrow t = 0.1973955598... \Rightarrow x \text{ or } a = \frac{1}{2} e^{-3(0.1973...)} \left( 4 - 3 \times \frac{12}{13} + 2 \sin(2 \times 0.1973...) \right) = ..$ <p>Finds a value of <math>t</math> from <math>\cos kt = c</math> (<math>k \neq 1, -1 &lt; c &lt; 1</math>) and uses their positive (or made positive) value of <math>t</math> to find a value of <math>x</math> (or <math>a</math>) via their PS. Accept a pair of stated values.</p> <p><b>Requires both previous M marks.</b></p>		ddM1
	$x \text{ or } a = 0.553(1164729...)$	awrt 0.553	A1
			(4)
			<b>Total 14</b>

Question Number	Scheme	Notes	Marks
7(a) Way 1	$w = \frac{z-3}{2i-z} \Rightarrow 2iw - wz = z-3 \Rightarrow z = \dots$	Attempts to make $z$ the subject and obtains any $f(w)$	M1
	$z = \frac{3+2iw}{w+1}$ or $\frac{-3-2iw}{-w-1}$	Any correct expression for $z$ in terms of $w$	A1
	$= \frac{3+2iu-2v}{u+iv+1} \times \frac{u+1-iv}{u+1-iv}$ Applies $w = u + iv$ and a correct multiplier for their $z$ seen or implied by a correct result from their $z$ . Denominator must have had a “ $w$ ”. Note alternative route below.		M1
	$x+iy = \frac{3+2iu-2v}{u+iv+1} \times \frac{u+1-iv}{u+1-iv} = \frac{(3-2v)(u+1)+2uv+2u(u+1)i-(3-2v)vi}{(u+1)^2+v^2}$ $y = x+3$ oe $\Rightarrow \frac{2u(u+1)-(3-2v)v}{(u+1)^2+v^2} = \frac{(3-2v)(u+1)+2uv}{(u+1)^2+v^2} + 3$ Multiplies, extracts real and imaginary parts and uses them in the equation $y = x + 3$ (oe) to produce an equation in $u$ and $v$ only – no “ $i$ ”s. Condone $y = \dots i$ if recovered. Can follow slips with multiplier but denominator of $z$ must have had a “ $w$ ” Note: Just $2u(u+1)-(3-2v)v = (3-2v)(u+1)+2uv+3$ is M0 (lost denominators)		M1
	$2u(u+1)-(3-2v)v = (3-2v)(u+1)+2uv+3(u+1)^2+3v^2$ $\Rightarrow u^2+7u+v^2+v+6=0$	Expands and simplifies to obtain an equation of a circle with 4 or 5 real unlike terms. <b>All previous Ms required.</b>	dddM1
<b>Alternative for the above 3 marks</b> (note this could be done by equating expressions for $y$ )			
$x+iy = \frac{3+2iu-2v}{u+iv+1} \Rightarrow (x+i(x+3))(u+1+iv) = 3+2ui-2v$ M1: Applies $z = x + iy$ , uses $y = x + 3$ and cross multiplies			
$x(u+1)-v(x+3)+(x+3)(u+1)i+xvi = 3-2v+2ui$ $\Rightarrow ux+x-vx-3v = 3-2v, \quad ux+x+3u+3+xv = 2u$ $\Rightarrow x = \frac{3+v}{u+1-v}, \quad x = \frac{-u-3}{u+1+v}$ M1: Equates real and imaginary parts and makes $x$ the subject twice			
$(3+v)(u+1+v) = -(u+3)(u+1-v) \Rightarrow 3u+3+3v+uv+v+v^2 = -u^2-u+uv-3u-3+3v$ $\Rightarrow u^2+v^2+7u+v+6=0$ M1: Equates expressions for $x$ to obtain a circle equation with 4 or 5 real unlike terms			
$\Rightarrow \left(u+\frac{7}{2}\right)^2 + \left(v+\frac{1}{2}\right)^2 = \frac{49}{4} + \frac{1}{4} - 6 = \frac{13}{2} \Rightarrow \text{centre: } \left(-\frac{7}{2}, -\frac{1}{2}\right) \text{ radius: } \frac{\sqrt{26}}{2} \text{ or } \sqrt{\frac{13}{2}}$ M1: Extracts the centre <b>and/or</b> radius from their <b>circle</b> equation, however obtained, with 4 or 5 real unlike terms. Circle equation must not be in terms of $z$ or $w$ . They must get one correct coordinate (but condone wrong sign) or the correct radius for their circle. May use $u^2+v^2+2gu+2fv+c=0 \Rightarrow \text{centre: } (-g, -f), \text{ radius} = \sqrt{g^2+f^2-c}$ A1: For a correct centre or radius <b>from a correct circle equation</b> A1: For correct centre and radius <b>from a correct circle equation</b> Centre as coordinates, $x/u = \dots, y/v = \dots$ or as $-\frac{7}{2} - \frac{1}{2}i$ and allow $(-\frac{7}{2}, -\frac{1}{2}i)$ Allow exact equivalents for coordinates/radius	M1 A1 A1		
			(8)



Question Number	Scheme	Notes	Marks
<b>7(a)</b>  <b>Way 2</b>	$w = \frac{z-3}{2i-z} = \frac{x+iy-3}{2i-x-iy} = \frac{x-3+i(x+3)}{2i-x-i(x+3)}$ [Note that it is possible to replace $x$ with $y-3$ ]	M1: Uses $z = x + iy$ and $y = x + 3$ in the given transformation A1: Correct expression for $w$ in terms of $x$	M1 A1
	$\frac{x-3+i(x+3)}{-x-i(x+1)} = u+iv \Rightarrow x-3+i(x+3) = -xu+iv(x+1) -iu(x+1) -ivx$	Applies $w = u + iv$ and multiplies	M1
	$x-3 = -ux+vx+v, \quad x+3 = -ux-u-ix$ $x = \frac{3+v}{1+u-v}, \quad x = \frac{-3-u}{1+u+v}$	Equates real and imaginary parts and makes $x$ the subject twice	M1
	$3+3u+3v+v+uv+v^2 = -3-3u+3v-u-u^2+uv$ $\Rightarrow u^2+v^2+7u+v+6=0$	Equates expressions for $x$ to obtain a circle equation with 4 or 5 real unlike terms. <b>All previous Ms required.</b>	dddM1
	$\Rightarrow \left(u+\frac{7}{2}\right)^2 + \left(v+\frac{1}{2}\right)^2 = \frac{49}{4} + \frac{1}{4} - 6 = \frac{13}{2} \Rightarrow \text{centre: } \left(-\frac{7}{2}, -\frac{1}{2}\right) \text{ radius: } \frac{\sqrt{26}}{2} \text{ or } \sqrt{\frac{13}{2}}$ M1: Applies a correct process to extract the centre and/or radius from a circle equation, however obtained, with 4 or 5 real unlike terms. One correct coordinate (but condone wrong sign) or radius correct for their circle. May use $u^2+v^2+2gu+2fv+c=0 \Rightarrow \text{centre: } (-g, -f), \text{ radius} = \sqrt{g^2+f^2-c}$ A1: For correct centre or radius from a correct circle equation A1: For correct centre and radius from a correct circle equation Centre as coordinates, $x/u=...$ , $y/v=...$ or as $-\frac{7}{2} - \frac{1}{2}i$ and allow $(-\frac{7}{2}, -\frac{1}{2}i)$ (8)	M1 A1 A1	
<b>Way 3</b>	e.g., 3 points on line are (0,3), (1,4) and (2,5) or $z_1 = 3i, z_2 = 1+4i, z_3 = 2+5i$	Attempts three points/complex numbers on $y = x + 3$ with 2 correct	M1
	$w = \frac{z-3}{2i-z} \Rightarrow w_1 = \frac{3i-3}{-i} \quad w_2 = \frac{-2+4i}{-1-2i} \quad w_3 = \frac{-1+5i}{-2-3i}$	Correct transformed complex numbers	A1
	$w_1 = \frac{3i-3}{-i} \times \frac{i}{i} \quad w_2 = \frac{-2+4i}{-1-2i} \times \frac{-1+2i}{-1+2i} \quad w_3 = \frac{-1+5i}{-2-3i} \times \frac{-2+3i}{-2+3i}$ At least two correct multipliers to remove "i" from denominator seen or implied (one if (-1, 2) used). Requires 2 correct points/complex numbers on line		M1
	$w_1 = -3-3i \quad w_2 = -\frac{6}{5} - \frac{8}{5}i \quad w_3 = -1-i$	Two correct complex numbers in $a + ib$ form or as points	M1
	e.g., $x^2 + y^2 + 2gx + 2fy + c = 0 \Rightarrow \frac{12}{5}g + \frac{16}{5}f - c = 0$ $6g+6f-c=18$ $2g+2f-c=0$	Uses a correct general equation of a circle to form three simultaneous equations. <b>All previous Ms required.</b>	dddM1
	$\Rightarrow g = \frac{7}{2}, f = \frac{1}{2}, c = 6 \Rightarrow \text{centre } (-g, -f): \left(-\frac{7}{2}, -\frac{1}{2}\right) \text{ radius} = \sqrt{g^2+f^2-c} = \frac{\sqrt{26}}{2} \text{ or } \sqrt{\frac{13}{2}}$ M1: Solves and obtains at least one correct coordinate (but condone wrong sign) or radius for their constants A1: Correct centre or radius from correct work A1: Correct centre and radius from correct work (8)	M1 A1 A1	

Question Number	Scheme	Notes	Marks
<b>7(b)</b> <b>(i) &amp; (ii)</b>		<p><b>M1: Any circle with the whole interior indicated.</b> Ignore any inconsistencies with their stated centre, value for radius (which may have been negative) or circle equation. If shaded, consider the shaded area but if not allow any credible indication such as an “<i>R</i>” inside the circle unless they have <b>clearly</b> indicated a segment.</p> <p><b>A1: Correct circle drawn in the correct position with whole interior shaded.</b> Entirely in quadrants 2 &amp; 3 and centre if marked in Q3 (if not marked then more than half of the circle in Q3). Condone if it appears that the area above the <i>x</i>-axis is greater than the area below provided the centre is indicated in Q3. <b>Must be shaded</b> but does not require a label. Circumference may be dotted/dashed line. Ignore incorrect labelling of centre/axes/intersections but <b>requires full marks in (a).</b></p>	<p><b>M1</b>  <b>(B1 on ePen)</b></p> <p><b>A1</b>  <b>(B1 on ePen)</b></p>
			(2)
			<b>Total 10</b>

Question Number	Scheme	Notes	Marks
<b>8(a)</b>	Allow “single fraction” to be implied by sum/difference of fractions with same denominator or a product of fractions. No further fractions in numerator/denominator.		

	$\cot 2x \{ + \tan x \} = \frac{\cos 2x}{\sin 2x} \left\{ + \frac{\sin x}{\cos x} \right\}$	Uses $\cot 2x = \frac{\cos 2x}{\sin 2x}$ or e.g., $\frac{\cos 2x}{2 \sin x \cos x}$	<b>M1</b>
	$\frac{\cos 2x + 2 \sin^2 x}{2 \sin x \cos x} \Rightarrow$ <p>e.g., <math>\frac{1 - 2 \sin^2 x + 2 \sin^2 x}{2 \sin x \cos x}</math> or <math>\frac{\cos^2 x - \sin^2 x + 2 \sin^2 x}{2 \sin x \cos x}</math></p> $\frac{2 \cos^2 x - 1 + 2 \sin^2 x}{\sin 2x}$ or $\frac{\cos 2x + 1 - \cos 2x}{\sin 2x}$	<p>Uses sufficient correct identities e.g.,</p> $\cos 2x = 1 - 2 \sin^2 x$ $\cos 2x = \cos^2 x - \sin^2 x$ $\cos 2x = 2 \cos^2 x - 1$ $2 \sin^2 x = 1 - \cos 2x$ $\cos 2x \cos x + \sin x \sin 2x = \cos(2x - x)$ <p>to obtain a correct single fraction with numerator in terms of <math>\sin x</math> and/or <math>\cos x</math> or “<math>\cos 2x + 1 - \cos 2x</math>”. A qualifying fraction must be seen before</p> $\frac{1}{2 \sin x \cos x} \text{ or } \frac{1}{\sin 2x}$ <p>Condone poor notation.</p>	<b>A1</b> <b>(M1 on ePen)</b>
	<p>OR <math>\frac{\cos 2x + \tan x \sin 2x}{\sin 2x} \Rightarrow</math></p> $\Rightarrow \frac{\cos 2x + \frac{\sin x}{\cos x} \times 2 \sin x \cos x}{\sin 2x} \Rightarrow \text{e.g., } \frac{1 - 2 \sin^2 x + 2 \sin^2 x}{\sin 2x}$ <p>OR <math>\frac{\cos 2x \cos x + \sin x \sin 2x}{\sin 2x \cos x} \Rightarrow</math></p> $\frac{\cos x}{\sin 2x \cos x} \text{ or } \frac{\cos^3 x - \sin^2 x \cos x + 2 \sin^2 x \cos x}{\sin 2x \cos x}$		
	$= \frac{1}{2 \sin x \cos x} \text{ or } \frac{1}{\sin 2x} = \operatorname{cosec} 2x^*$	Fully correct proof with one of the two intermediate fractions seen. All notation correct – no mixed or missing arguments or e.g. $\sin x^2$ for this mark.	<b>A1*</b>
<b>(3)</b>			
<b>Alt</b>	$\cot 2x \{ + \tan x \} = \frac{1 - \tan^2 x}{2 \tan x} \{ + \tan x \}$	Uses $\cot 2x = \frac{1 - \tan^2 x}{2 \tan x}$	<b>M1</b>
	$\frac{1 - \tan^2 x + 2 \tan^2 x}{2 \tan x} \Rightarrow$ <p>e.g., <math>\frac{\tan^2 x + 1}{2 \tan x} \Rightarrow \frac{\left(\frac{\sin x}{\cos x}\right)^2 + 1}{2 \frac{\sin x}{\cos x}} \Rightarrow \frac{\cos x (\sin^2 x + \cos^2 x)}{2 \cos^2 x \sin x}</math></p> <p>or <math>\frac{\tan^2 x + 1}{2 \tan x} \left\{ \times \frac{\cos x}{\cos x} \right\} \Rightarrow \frac{\sin^2 x + \cos^2 x}{2 \sin x \cos x}</math></p> <p>or <math>\frac{\sec^2 x}{2 \tan x}</math> or <math>\frac{\cos x}{2 \cos^2 x \sin x}</math></p>	<p>Uses correct identities e.g.,</p> $\tan x = \frac{\sin x}{\cos x} \text{ oe}$ <p>to obtain a correct single fraction in <math>\sin x</math> and <math>\cos x</math> but allow <math>\frac{\sec^2 x}{2 \tan x}</math> following use of</p> $\sec^2 x = 1 + \tan^2 x$ <p>A qualifying fraction must be seen before</p> $\frac{1}{2 \sin x \cos x} \text{ or } \frac{1}{\sin 2x}$ <p>Condone poor notation.</p>	<b>A1</b> <b>(M1 on ePen)</b>
	$\frac{1}{2 \sin x \cos x} \text{ or } \frac{1}{\sin 2x} = \operatorname{cosec} 2x^*$	Fully correct proof with one of the two intermediate fractions seen. All notation correct – no mixed or missing arguments or e.g. $\sin x^2$ for this mark.	<b>A1*</b>
<b>(3)</b>			

Question Number	Scheme	Notes	Marks
<b>8(b)</b>	Examples:		<b>M1</b> <b>A1</b>

	$y^2 = w \sin 2x \Rightarrow 2y \frac{dy}{dx} = \frac{dw}{dx} \sin 2x + 2w \cos 2x$ $\text{or } y = w^{\frac{1}{2}} (\sin 2x)^{\frac{1}{2}} \Rightarrow \frac{dy}{dx} = \frac{1}{2} w^{-\frac{1}{2}} (\sin 2x)^{-\frac{1}{2}} (2 \cos 2x) + \frac{1}{2} w^{-\frac{1}{2}} \frac{dw}{dx} (\sin 2x)^{\frac{1}{2}}$ $\text{or } w = \frac{y^2}{\sin 2x} \Rightarrow \frac{dw}{dx} = \frac{2y \sin 2x \frac{dy}{dx} - y^2 \cdot 2 \cos 2x}{\sin^2 2x}$ $\text{or } w = y^2 \operatorname{cosec} 2x \Rightarrow 2y \frac{dy}{dx} \operatorname{cosec} 2x - 2y^2 \operatorname{cosec} 2x \cot 2x$ <p><b>M1:</b> Attempts the differentiation of the given substitution using the product/quotient and chain rules <b>and</b> obtains an <b>equation</b> in <math>\frac{dy}{dx}</math> and <math>\frac{dw}{dx}</math> of the correct form (sign/coefficient errors only and allow sign errors with quotient/product rule).</p> <p>This mark is not available for work in <math>\frac{dy}{dw}</math> or <math>\frac{dw}{dy}</math> unless appropriate work follows to achieve an equation in <math>\frac{dy}{dx}</math> and <math>\frac{dw}{dx}</math> of the correct form.</p> <p><b>A1:</b> Correct differentiation</p>	
	$y \frac{dy}{dx} + y^2 \tan x = \sin x \rightarrow \text{e.g., } \frac{1}{2} \left( \frac{dw}{dx} \sin 2x + 2w \cos 2x \right) + w \sin 2x \tan x = \sin x$ <p>A recognisable attempt to eliminate <math>y</math> from the original equation to obtain an equation involving <math>\frac{dw}{dx}</math>, <math>w</math> and <math>x</math> only. Not dependent.</p>	<b>M1</b>
	$\Rightarrow \frac{dw}{dx} + 2w (\cot 2x + \tan x) = \frac{2 \sin x}{\sin 2x}$ $\Rightarrow \frac{dw}{dx} + 2w \operatorname{cosec} 2x = \sec x *$ <p>Fully correct work leading to the given equation with <math>2w (\cot 2x + \tan x)</math> or e.g., <math>2w \cot 2x + 2w \tan x</math> clearly replaced by <math>2w \operatorname{cosec} 2x</math> but allow <math>\cot 2x</math> written as <math>\frac{1}{\tan 2x}</math> or <math>\frac{\cos 2x}{\sin 2x}</math> and/or <math>\tan x</math> written as <math>\frac{\sin x}{\cos x}</math></p> <p>If the result in (a) is not clearly used there must be full equivalent work. Allow use of "csc <math>2x</math>"</p>	<b>A1*</b>
		<b>(4)</b>

Question Number	Scheme	Notes	Marks
-----------------	--------	-------	-------

<p><b>8(c)</b></p> $\frac{dw}{dx} + 2w \operatorname{cosec} 2x = \sec x \Rightarrow \text{IF} = e^{2 \int \operatorname{cosec} 2x dx} = \tan x$ <p>or <math>e^{-\ln(\operatorname{cosec} 2x + \cot 2x)} \Rightarrow \frac{1}{\operatorname{cosec} 2x + \cot 2x}</math> or <math>\frac{1}{\cot x}</math> or <math>\tan x</math></p>		<p><b>M1:</b> <math>e^{2 \int \operatorname{cosec} 2x (dx)}</math> condoning omission of one or both “2”s</p> <p><b>A1:</b> <math>\tan x</math> oe</p> <p>Allow <math>k \tan x</math> e.g., <math>e^{2c} \tan x</math></p> <p>Not just <math>e^{\ln(\tan x)}</math></p>	<p><b>M1</b> <b>A1</b></p>
	$\Rightarrow w \tan x = \int \tan x \sec x \{dx\}$	<p>Correctly applies their integrating factor to the equation, i.e.,</p> $\Rightarrow \text{IF} \times w = \int \text{IF} \times \sec x \{dx\}$ <p>Allow equivalents for <math>\sec x</math>. Condone “y” used for “w” for this mark.</p>	<p><b>M1</b></p>
	$\Rightarrow w \tan x = \sec x (+c)$	<p>Correct equation oe with or without constant.</p>	<p><b>A1</b></p>
	<p>Using IF = <math>\frac{1}{\operatorname{cosec} 2x + \cot 2x} \Rightarrow</math> RHS of <math>\int \frac{\sec x}{\operatorname{cosec} 2x + \cot 2x} dx</math> which is likely to need rewriting as <math>\int \tan x \sec x dx</math></p> <p>Note that IBP on <math>\sec x \tan x</math> by writing it as <math>\sec^2 x \sin x</math> can lead to <math>\sin x \tan x + \cos x (+c)</math></p> <p>Use Review for any attempts at integration you are unsure about.</p>		
	<p>e.g., <math>y^2 = w \sin 2x</math> and <math>w \tan x = \sec x + c \Rightarrow \frac{y^2}{\sin 2x} \tan x = \sec x + c</math></p> $\Rightarrow y^2 = \dots \left\{ \frac{\sin 2x}{\tan x} (\sec x + c) \right\}$ <p>Substitutes for <math>w</math> correctly and reaches <math>y^2 = \dots</math></p> <p>Their <math>y^2 = \dots</math> must be consistent with <b>their</b> equation in <math>w</math> and <math>x</math> that immediately followed their integration.</p> <p><b>This mark requires both previous M marks and an attempt at integration that includes a “+ c”</b></p> <p>A further example is:</p> $w = \operatorname{cosec} x + \frac{c}{\tan x} \Rightarrow y^2 = \operatorname{cosec} x \sin 2x + \frac{c \sin 2x}{\tan x}$		<p><b>ddM1</b></p>
	$\left\{ \text{e.g., } y^2 = \frac{2 \sin x \cos^2 x}{\sin x} \left( \frac{1}{\cos x} + c \right) \Rightarrow \right\}$ $y^2 = 2 \cos x + A \cos^2 x$ <p>Any correct <math>y^2 = \dots</math> equation with RHS fully in terms of <math>\cos x</math>. E.g. accept</p> $y^2 = 2 \cos x + 2c \cos^2 x \quad y^2 = \cos x (2 + A \cos x) \quad y^2 = 2 \cos^2 x \left( \frac{1}{\cos x} + c \right)$ <p>Ignore any inconsistencies with the constant e.g., <math>2c</math> later written as <math>c</math></p>		<p><b>A1</b></p>
			<p><b>(6)</b></p>
			<p><b>Total 13</b></p>

