

**OXFORD**

INTERNATIONAL  
AQA EXAMINATIONS

---

# INTERNATIONAL A-LEVEL MATHEMATICS

## MA04

(9660/MA04) Unit S2 Statistics

---

Mark scheme

January 2023

---

Version: 1.0 Final



2 3 1 X M A 0 4 / M S

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from [oxfordaqaexams.org.uk](http://oxfordaqaexams.org.uk)

#### **Copyright information**

OxfordAQA retains the copyright on all its publications. However, registered schools/colleges for OxfordAQA are permitted to copy material from this booklet for their own internal use, with the following important exception: OxfordAQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2023 Oxford International AQA Examinations and its licensors. All rights reserved.

**Key to mark scheme abbreviations**

<b>M</b>	Mark is for method
<b>m</b>	Mark is dependent on one or more M marks and is for method
<b>A</b>	Mark is dependent on M or m marks and is for accuracy
<b>B</b>	Mark is independent of M or m marks and is for method and accuracy
<b>E</b>	Mark is for explanation
<b>√ or ft</b>	Follow through from previous incorrect result
<b>CAO</b>	Correct answer only
<b>CSO</b>	Correct solution only
<b>AWFW</b>	Anything which falls within
<b>AWRT</b>	Anything which rounds to
<b>ACF</b>	Any correct form
<b>AG</b>	Answer given
<b>SC</b>	Special case
<b>oe</b>	Or equivalent
<b>A2, 1</b>	2 or 1 (or 0) accuracy marks
<b>-x EE</b>	Deduct x marks for each error
<b>NMS</b>	No method shown
<b>PI</b>	Possibly implied
<b>SCA</b>	Substantially correct approach
<b>sf</b>	Significant figure(s)
<b>dp</b>	Decimal place(s)

Q	Answer	Marks	Comments
1(a)	$E(3Y - 2X) = 3 \times 15 - 2 \times 12$	M1	
	$= 21$	A1	
		2	

Q	Answer	Marks	Comments
1(b)	$\text{Var}(3Y - 2X) = 3^2 \times 2.5 + 2^2 \times 5$	M1	Condone a sign error
	$= 42.5$	A1	oe eg $85/2$ $85 \div 2$ $\frac{85}{2}$
		2	

	<b>Question 1 Total</b>	<b>4</b>	
--	-------------------------	----------	--

Q	Answer	Marks	Comments
<b>2(a)</b>	Stages 1, 4 and 8 contain errors	<b>B1</b>	At least two stages correctly identified
	Stage 1 – Hypothesis are wrong should be $H_0: \mu = 18.3$ $H_1: \mu \neq 18.3$	<b>E1</b>	Clear comment regarding population parameter $\mu$ not the sample $\bar{x}$
	Stage 4 – [ $z_{\text{critical}} = 1.2816$ ] should be $z_{\text{critical}} = (\pm)1.6449$	<b>E1</b>	Allow $\pm$ but not –
	Stage 8 – wrong conclusion: should not reject $H_0$ : Insufficient evidence to suggest a change in mean has occurred at the 10% level of significance	<b>E1</b>	<b>oe</b>
		<b>4</b>	

Q	Answer	Marks	Comments
<b>2(b)</b>	The sample was taken from a normal distribution so the sample is also normally distributed.	<b>E1</b>	<b>oe</b> for example 'a sample inherits the characteristics of the parent population'
		<b>1</b>	

Q	Answer	Marks	Comments
<b>2(c)</b>	A Critical Region	<b>B1</b>	
		<b>1</b>	

	<b>Question 2 Total</b>	<b>6</b>	
--	-------------------------	----------	--

Q	Answer	Marks	Comments
3(a)	$\frac{dF(t)}{dt} = \frac{3t^2}{64}$	M1	Obtains $kt^2$
	$f(t) = \frac{3t^2}{64}, \quad 0 \leq t \leq 4$	A1	Requires both parts
	0, otherwise	B1	
		3	

Q	Answer	Marks	Comments
3(b)(i)	$[\mu = E(T) =] \int_0^4 t \times \frac{3t^2}{64} dt = \left[ \frac{3t^4}{256} \right]_0^4$	M1	Attempt to integrate their $tf(t)$ with correct limits PI
	= 3	A1	
		2	

Q	Answer	Marks	Comments
3(b)(ii)	$[E(T^2) =] \int_0^4 t^2 \times \frac{3t^2}{64} dt = \left[ \frac{3t^5}{320} \right]_0^4$	M1	Attempt to integrate their $t^2f(t)$ with correct limits PI
	= 9.6	A1ft	PI, oe
	$\sigma^2 = [E(T^2) - E(T)^2 =] 9.6 - 3^2$	M1	ft their $E(t^2) - E(t)^2$ PI
	$\sigma = \sqrt{0.6} = 0.775 \text{ (3 sf)}$	A1	Condone $\sigma = 0.6$ CAO to 3 sf
		4	

Q	Answer	Marks	Comments
3(c)	$P(3 - 2\sqrt{0.6} \leq T \leq 3 + 2\sqrt{0.6})$ $= P(1.45[0806662] \leq T \leq 4.54[9193338])$ $F(4.54) - F(1.45[0806662])$ $F(4.54) = [F(4) = ] 1$ $1 - 0.04771431013 = 0.952 \text{ (3 sf)}$	<p><b>M1</b></p> <p><b>M1</b></p> <p><b>B1</b></p> <p><b>A1</b></p>	<p>Allow their <math>\mu</math> &amp; <math>\sigma</math> <b>PI</b></p> <p><b>PI</b> by correct answer</p> <p><b>AWRT</b> 0.952</p>
		<b>4</b>	
	<b>Question 3 Total</b>	<b>13</b>	

Q	Answer	Marks	Comments
4	$H_0: \mu = 25$ $H_1: \mu < 25$ $\bar{x} = 24.86$ $s^2 = \frac{1}{500-1} \times \left( 310000 - \frac{12430^2}{500} \right)$ $= 1.98 [4368737...]$ $\bar{X} \sim N \left( 25, \frac{1.98 [...]}{500} \right)$ $z = \frac{24.86 - 25}{\sqrt{\frac{1.98 [...]}{500}}}$ $= -2.2 [22295719...]$ $z_{\text{critical}} = -2.3263$ <p>Do not reject <math>H_0</math> as  <math>-2.2 [222...] &gt; -2.3 [263...]</math> or <math>z &gt; z_{\text{crit}}</math></p> <p>Evidence to suggest that the hand-drying time does not have a mean lower than 25 [at the 1% level of significance]</p>	<p><b>B1</b></p> <p><b>B1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>B1</b></p> <p><b>A1ft</b></p> <p><b>E1ft</b></p>	<p>Both hypotheses</p> <p>Attempt at variance formula                      Allow one slip                      Implied by correct answer</p> <p><b>AWRT</b> 1.98                      Accept <math>s = 1.40 [8676236]</math></p> <p><b>PI</b>, <math>\bar{X} \sim N \left( 25, \frac{s^2}{500} \right)</math></p> <p>Calculates <math>z</math> with their <math>s^2</math></p> <p><b>AWRT</b> <math>-2.2</math> or <math> z  = 2.2</math>                      or probability of 0.013</p> <p><b>AWRT</b> <math>-2.3</math>                      Allow <math>z_{\text{critical}} = \pm 2.3263</math></p> <p><b>ft</b> their <math>z</math> and <math>z_{\text{crit}}</math> provided signs are consistent</p> <p>Allow a correct comparison of <math>z_{\text{crit}}</math> and <math> z </math>                      or compares their 0.013 to 0.01 and correct statement</p> <p>Must be consistent with their conclusion on whether or not to reject <math>H_0</math>, or their <math>z</math> and <math>z_{\text{crit}}</math> if not explicitly stated</p> <p>Condone definite statement</p>
		<b>10</b>	
	<b>Question 4 Total</b>	<b>10</b>	



Q	Answer	Marks	Comments
5(a)(i)	Condition 1: The number of events/trials, $n$ must be large; and  Condition 2: The probability of a successful outcome, $p$ , must be small	B1	$n$ must be large <b>and</b> the $p$ value must be small  Allow <b>eg</b> $n \geq 50$ and $p \leq 0.1$
		1	

Q	Answer	Marks	Comments
5(a)(ii)	[Mean = $np$ = ] $\lambda = 500 \times 0.02 = 10$	B1	
		1	

Q	Answer	Marks	Comments
5(a)(iii)	$P(X < 10) = P(X \leq 9)$  [ = 0.4579297391... ]  = 0.458	M1  A1	PI  AWRT 0.458
		2	

Q	Answer	Marks	Comments
5(b)(i)	$H_0: \lambda = 8$ $H_1: \lambda \neq 8$  $X = 5$ [per hectare]  $P(X \leq 5) = 0.191$ [2360621...]  $0.191 > 0.05$  Do not reject $H_0$  Insufficient evidence to suggest the mean number of ground nests [per hectare] has changed from 8 [at the 10% level of significance]	B1  M1  A1  A1  B1ft  E1	Allow $H_0: \lambda = 240$ $H_1: \lambda \neq 240$  PI  Comparison of their probabilities or a lower CR of $X \leq 3$  Correct conclusion on $H_0$ based on their comparison of test statistic with critical value  Must be in context, must not be definite and all the previous 5 marks must have been awarded
		6	

Q	Answer	Marks	Comments
5(b)(ii)	$P(X \leq c) < 0.05$ $P(X \leq 3) = 0.0424$ [or $P(X \geq 4) = 0.9576$ ] and $P(X \leq 4) = 0.0996$ [or $P(X \geq 5) = 0.9004$ ]	M1	PI by correct CR, could be written as a single CR later
	$P(X \geq c) < 0.05$ $P(X \geq 14) = 0.0342$ [or $P(X \leq 13) = 0.9658$ ] and $P(X \geq 13) = 0.0638$ [or $P(X \leq 12) = 0.9362$ ]	M1	PI by correct CR, could be written as a single CR later
	Both included CR = $\{0, 1, 2, 3, 14, 15, \dots\}$	A1	Allow $X \leq 3$ and $X \geq 14$ oe
		3	

Q	Answer	Marks	Comments
5(c)(i)	Concluding that the mean number of ground nests [per hectare] has changed [from 8] even though it hasn't.	B1	
		1	

Q	Answer	Marks	Comments
5(c)(ii)	$[0.0424 + 0.0342 =] 0.0766$	B1	CAO
		1	

	<b>Question 5 Total</b>	<b>15</b>	
--	-------------------------	-----------	--

Q	Answer	Marks	Comments
6(a)	$P(X_1 < 10) = P\left(Z < \frac{10 - 12}{1.5}\right)$ $= 0.0912[1121973\dots]$	M1	Attempt to standardise <b>PI</b> <b>oe, e.g.</b> $P(Z < -1.33[3\dots])$ or $1 - P(Z < 1.33[3\dots])$
		A1	<b>AWFW</b> 0.0912 to 0.0918
		<b>2</b>	

Q	Answer	Marks	Comments
6(b)	$P(X_2 < a) = 0.9$ $z = 1.28[1551638\dots]$ $\frac{a - 8}{0.8} = 1.28$ $9.03 \text{ [mins] (to 3 sf)}$	B1	<b>Allow</b> $\pm 1.28[1551638\dots]$
		M1	Forms an equation with their $1.2 <  z  < 1.3$
		A1	9.025[241311\dots] <b>AWRT</b> 9.03 <b>oe, e.g.</b> 9 minutes, 2 seconds
			<b>3</b>

Q	Answer	Marks	Comments
6(c)	$z = 0.915[3645453]$ or $z = [-]0.075[26985986]$ $0.915[3645453] = \frac{16 - \mu}{\sigma}$ or $-0.075[26985986] = \frac{10 - \mu}{\sigma}$ $0.915[3645453] = \frac{16 - \mu}{\sigma}$ and $-0.075[26985986] = \frac{10 - \mu}{\sigma}$ $(0.915[3645453] + 0.075[26985986])\sigma$ $\qquad\qquad\qquad = 16 - 10$ $\sigma = 6.06$ (3 sf) $\mu = 10.5$ (3 sf)	B1  M1  A1  M1  A1  A1	PI by correct standardisation 0.9154 or 0.0753 from tables  oe At least one correct standardisation PI by at least one correct value for $\mu$ or $\sigma$  oe Both correct standardisations PI by at least one correct value for $\mu$ or $\sigma$ Attempt to eliminate one variable simultaneously PI by at least one correct value for $\mu$ or $\sigma$  AFWW 6.05 to 6.06  AWRW 10.5
		6	

Q	Answer	Marks	Comments
6(d)	$\mu - 2\sigma < 0$ or $\mu - 3\sigma < 0$ or time cannot be less than zero in this context	B1	A clear comment about the lower tail having $t < 0$
		1	

Q	Answer	Marks	Comments
6(e)	$T = X_1 + X_2 + X_3 + X_4$ $T \sim N(12 + 8 + 7 + 14, 1.5^2 + 0.8^2 + 1^2 + 2^2)$ $T \sim N(41, 7.89)$ $P(T < 35) = P\left(Z < \frac{35 - 41}{\sqrt{7.89}}\right)$ $[= P(Z < -2.136056564...)]$ $= 0.0163[3740189...]$	M1  M1  A1	PI  AFWW 0.0162 to 0.0164
		3	

	<b>Question 6 Total</b>	<b>15</b>	
--	-------------------------	-----------	--

Q	Answer	Marks	Comments
7(a)	$P(X=4) = \frac{e^{-2.5} \times 2.5^4}{4!}$ $= 0.1336[018858]$	M1	PI
		A1	
		2	

Q	Answer	Marks	Comments
7(b)	$\lambda = 6 \times 2.5 = 15$ $P(X > 18) = 1 - P(X \leq 18)$ $= 1 - 0.8194[717351]$ $= 0.1805 \text{ (to 4 sf)}$	B1	PI
		M1	
		A1	AWRT 0.1805
		3	

Q	Answer	Marks	Comments
7(c)(i)	$\left[ \frac{1}{\lambda} = \right] 0.4$	B1	
		1	

Q	Answer	Marks	Comments
7(c)(ii)	$\left[ \left( \frac{1}{\lambda} \right)^2 = 0.4^2 = \right] 0.16$	B1	
		1	

Q	Answer	Marks	Comments
7(d)	$1 - e^{-2.5c} = 0.9$ $e^{-2.5c} = 0.1$ $-2.5c = \ln 0.1$ $c = \frac{2}{5} \ln 10$ $c = 0.9210[340372\dots] \text{ [weeks]}$ $c = 6 \text{ [days]}$	M1	$1 - e^{-kc} = 0.9$ for $k > 0$ or correct integration with correct limits
		M1	
		A1	Finds their $c$ in exact form or as a value for the number of weeks
		3	

	<b>Question 7 Total</b>	<b>10</b>	
--	-------------------------	-----------	--

Q	Answer	Marks	Comments
8(a)	$\left[ F(x) = \int_2^x f(x) dx = \right] \int_2^x \frac{96}{(5x+k)^2} dx$ $= \left[ \frac{96}{-5(5x+k)} \right]_2^x$ $= -\frac{96}{5} \left( \frac{1}{5x+k} - \frac{1}{10+k} \right)$ $= -\frac{96}{5} \left( \frac{(10+k) - (5x+k)}{(10+k)(5x+k)} \right)$ <p>or</p> $= \frac{96}{5} \left( \frac{5x-10}{(10+k)(5x+k)} \right)$ <p>or</p> $= \frac{96}{5} \left( \frac{5(x-2)}{(10+k)(5x+k)} \right)$ $F(x) = \begin{cases} 0 & x < 2 \\ \frac{96(x-2)}{(10+k)(5x+k)} & 2 \leq x \leq d \\ 1 & x > d \end{cases}$	<p><b>M1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>A1</b></p>	<p>Attempt at integration to obtain <math>a(5x+k)^{-1}</math> with <math>a \neq 96</math></p> <p><b>oe</b> Substitute limits into their integration of the form <math>a(5x+k)^{-1}</math> with <math>a \neq 96</math></p> <p><b>oe</b> Correct single fraction</p> <p><b>AG</b> Must be convincingly shown</p> <p><b>SC2</b> for correctly differentiating <math>F(x)</math> w.r.t. to <math>x</math> to show <math>f(x) = \frac{96}{(5x+k)^2}</math></p>
		<b>4</b>	

Q	Answer	Marks	Comments
8(b)	$[F(4.4) = 0.8 \Rightarrow] 0.8 = \frac{96(4.4 - 2)}{(10 + k)(5 \times 4.4 + k)}$ $220 + 32k + k^2 = 288$ $k^2 + 32k - 68 = 0$ $(k - 2)(k + 34) = 0$ $k = 2$	<p><b>M1</b></p> <p><b>m1</b></p> <p><b>A1</b></p>	<p><b>oe</b></p> <p><math>0.8k^2 + 25.6k - 54.4 = 0</math>                      Attempts to solve the correct quadratic equation  <b>PI</b> by <math>k = 2</math></p> <p>Must not include <math>k = -34</math></p>
		<b>3</b>	
	<b>Question 8 Total</b>	<b>7</b>	