

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Science-Specific Marking Principles

- 1 Examiners should consider the context and scientific use of any keywords when awarding marks. Although keywords may be present, marks should not be awarded if the keywords are used incorrectly.
- 2 The examiner should not choose between contradictory statements given in the same question part, and credit should not be awarded for any correct statement that is contradicted within the same question part. Wrong science that is irrelevant to the question should be ignored.
- 3 Although spellings do not have to be correct, spellings of syllabus terms must allow for clear and unambiguous separation from other syllabus terms with which they may be confused (e.g. ethane / ethene, glucagon / glycogen, refraction / reflection).
- 4 The error carried forward (ecf) principle should be applied, where appropriate. If an incorrect answer is subsequently used in a scientifically correct way, the candidate should be awarded these subsequent marking points. Further guidance will be included in the mark scheme where necessary and any exceptions to this general principle will be noted.
- 5 'List rule' guidance
For questions that require *n* responses (e.g. State **two** reasons ...):
 - The response should be read as continuous prose, even when numbered answer spaces are provided.
 - Any response marked *ignore* in the mark scheme should not count towards *n*.
 - Incorrect responses should not be awarded credit but will still count towards *n*.
 - Read the entire response to check for any responses that contradict those that would otherwise be credited. Credit should **not** be awarded for any responses that are contradicted within the rest of the response. Where two responses contradict one another, this should be treated as a single incorrect response.
 - Non-contradictory responses after the first *n* responses may be ignored even if they include incorrect science.

6 Calculation specific guidance

Correct answers to calculations should be given full credit even if there is no working or incorrect working, **unless** the question states 'show your working'.

For questions in which the number of significant figures required is not stated, credit should be awarded for correct answers when rounded by the examiner to the number of significant figures given in the mark scheme. This may not apply to measured values.

For answers given in standard form (e.g. $a \times 10^n$) in which the convention of restricting the value of the coefficient (a) to a value between 1 and 10 is not followed, credit may still be awarded if the answer can be converted to the answer given in the mark scheme.

Unless a separate mark is given for a unit, a missing or incorrect unit will normally mean that the final calculation mark is not awarded. Exceptions to this general principle will be noted in the mark scheme.

7 Guidance for chemical equations

Multiples / fractions of coefficients used in chemical equations are acceptable unless stated otherwise in the mark scheme.

State symbols given in an equation should be ignored unless asked for in the question or stated otherwise in the mark scheme.

Question	Answer	Marks
1(a)(i)	Value of L with unit and in the range 20.0–25.0 cm.	1
1(a)(ii)	Value of T in range 0.80–1.20 s.	1
	Repeats: at least two measurements of at least $5T$.	1
1(b)	Six sets of readings of L and T with correct trend and without help from the Supervisor scores 4 marks, five sets scores 3 marks, etc.	4
	Range: $L_{\min} \leq 12.0$ cm and $L_{\max} \geq 40.0$ cm.	1
	Column headings: Each column heading must contain a quantity and a unit where appropriate. The presentation of quantity and unit must conform to accepted scientific convention e.g. L^2 / cm^2 .	1
	Consistency: All values of raw L must be given to the nearest mm.	1
	Significant figures: Values of L^2 should be to the same number of s.f. as (or one more than) the number of s.f. in the corresponding value of L .	1
	Calculation: Values of L^2 calculated correctly.	1

PUBLISHED

Question	Answer	Marks
1(c)(i)	<p>Axes: Sensible scales must be used, no awkward scales (e.g. 3:10 or fractions). Scales must be chosen so that the plotted points occupy at least half the graph grid in both the x and y directions. Axes must be labelled with the quantity that is being plotted. Scale markings should be no more than three large squares apart.</p>	1
	<p>Plotting of points: All observations in the table must be plotted on the grid. Diameter of plotted points must be \leq half a small square. Points must be plotted to an accuracy of half a small square in both x and y directions.</p>	1
	<p>Quality: All points in the table (at least 5) must be plotted on the grid. Trend of points on graph must be correct. It must be possible to draw a straight line that is within $\pm 0.02 \text{ m}^2$ on the L^2 axis (x-axis) of <u>all</u> plotted points.</p>	1
1(c)(ii)	<p>Line of best fit: Judge by the balance of all points on the grid about the candidate's line (at least 5 points). There must be an even distribution of points either side of the line along the full length. Allow one anomalous point only if clearly indicated (i.e. circled or labelled) by the candidate. There must be at least five points left after the anomalous point is disregarded. Line must not be kinked or thicker than half a small square.</p>	1
1(c)(iii)	<p>Gradient: The hypotenuse of the triangle used must be greater than half the length of the drawn line. Both read-offs must be accurate to half a small square in both the x and y directions. Method of calculation must be correct, e.g. $\Delta y / \Delta x$. Gradient sign on answer line matches graph drawn.</p>	1
	<p>y-intercept: Correct read-off from a point on the line and substituted into $y = mx + c$ or an equivalent expression. Read-off accurate to half a small square in both x and y directions. or Intercept read directly from the graph, with read-off at $L^2 = 0$, accurate to half a small square.</p>	1

Question	Answer	Marks
1(d)	a equal to candidate's gradient and b equal to candidate's intercept. Values must not be written as fractions.	1
	Units for a (e.g. s^2cm^{-2}) and b (e.g. s^2) are correct.	1

COURSEWORK

PUBLISHED

Question	Answer	Marks
2(a)(i)	Evidence of measuring a multiple of t and then dividing.	1
2(a)(ii)	Values of d_1 and d_2 to nearest 0.1 cm.	1
2(a)(iii)	Correct calculation of V_R .	1
2(a)(iv)	Justification based on significant figures in t , d_1 and d_2 .	1
2(b)(i)	Value of x_1 to nearest 1 cm ³ and in range 45–55 cm ³ .	1
2(b)(ii)	Value of x_2 less than x_1 .	1
	Correct calculation of V_A .	1
2(b)(iii)	Percentage uncertainty based on an absolute uncertainty in the range 2–4 cm ³ . If repeated readings have been taken, then the uncertainty can be half the range (but not zero) if working is clearly shown. Correct method of calculation to find percentage uncertainty.	1
2(c)	Second values for x_1 and x_2 .	1
	Second $V_A >$ first V_A .	1
2(d)(i)	Two values of k calculated correctly. The final values must not be written as fractions.	1
2(d)(ii)	Valid comment consistent with the calculated values of k , testing against a criterion specified by the candidate.	1

PUBLISHED

Question	Answer	Marks
2(e)(i)	<p>A Two readings are not enough to draw a (valid) conclusion (not “not enough for accurate results”, “few readings”).</p> <p>B Large percentage uncertainty in $t/d_1/d_2/V_R$.</p> <p>C Difficult to remove all air from cup at start/air leaks out of cup when operating syringe.</p> <p>D Difficult to judge when cup starts to rise or difficult to operate plunger smoothly or difficult to stop plunger when cup starts to rise.</p> <p>E Cup sticks to wall of container.</p> <p>F <u>Volume (or mass)</u> of cup/paper clip/string not taken into account.</p> <p>G <u>x (or V_A)</u> values affected by water getting into tube.</p> <p><i>1 mark for each point up to a maximum of 4.</i></p>	4

PUBLISHED

Question	Answer	Marks
2(e)(ii)	<p>A Take more readings <u>and</u> plot a graph or take more readings <u>and</u> compare k values (not “repeat readings” on its own).</p> <p>B Use vernier calipers/digital calipers/micrometer/travelling microscope.</p> <p>C Description of workable method of removing air.</p> <p>D Video/film/record with <u>syringe</u> in view or mark cup starting position on container.</p> <p>E Use wider container.</p> <p>F Method of finding volume of cup/string/paper clip or method of measuring mass of cup/string/paper clip e.g. top pan balance.</p> <p>G Method of removing water from tube or use new tube.</p> <p><i>1 mark for each point up to a maximum of 4.</i></p>	4